Full wwPDB NMR Structure Validation Report (i)

May 7, 2024 – 12:16 pm BST

PDB ID	:	5A4H
BMRB ID	:	25684
Title	:	Solution structure of the lipid droplet anchoring peptide of CGI-58 bound to DPC micelles
Authors	:	Boeszoermenyi, A.; Arthanari, H.; Wagner, G.; Nagy, H.M.; Zangger, K.; Lindermuth, H.; Oberer, M.
Deposited on	:	2015-06-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 46%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	39	8% 5% 13% 74%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 16 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:18-A:27 (10)	0.09	16

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20
2	1, 3, 4, 6, 7
3	2, 5
Single-model clusters	8

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 531 atoms, of which 253 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	39	531	172	253	46	58	2	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	GLY	-	expression tag	UNP Q9DBL9
A	6	ALA	-	expression tag	UNP Q9DBL9
A	7	MET	-	expression tag	UNP Q9DBL9
A	8	GLY	-	expression tag	UNP Q9DBL9
A	9	SER	-	expression tag	UNP Q9DBL9

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%

G5 A6 M7 G8 S9 V10 D11 S12 A13 D14 A15 G16 G17 G18 S19 G20 W21 L22 T23 G24 W25 L26 P27 T28 W29 C30 P31 T32 S33 T34 S35 H36 L37 K38 E39 A40 E41 E42 K43

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%

G5 A6 M7 G8 S9 V10 D11 S12 A13 D14 A15 G16 G17 G18 S19 G20 W21 L22 T23 G24 W25 L26 P27 T28 W29 C30 P31 T32 S33 T34 S35 H36 L37 K38 E39 A40 E41 E42 K43

4.2.2 Score per residue for model 2

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 10% • 13% 74%

G5 A6 M7 G8 S9 V10 D11 S12 A13 D14 A15 G16 G17 G20 W21 L22 T23 G24 W25 L26 P27 T28 W29 C30 P31 T32 S33 T34 S35 H36 L37 K38 E39 A40 E41 E42 K43

4.2.3 Score per residue for model 3

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%



4.2.4 Score per residue for model 4

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



4.2.5 Score per residue for model 5

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%



4.2.6 Score per residue for model 6

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



4.2.7 Score per residue for model 7

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



4.2.8 Score per residue for model 8

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 8% 10% 74%



4.2.9 Score per residue for model 9

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



4.2.10 Score per residue for model 10

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%



4.2.11 Score per residue for model 11

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%



4.2.12 Score per residue for model 12

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%



4.2.13 Score per residue for model 13

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%



4.2.14 Score per residue for model 14

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



4.2.15 Score per residue for model 15

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



4.2.16 Score per residue for model 16 (medoid)

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



4.2.17 Score per residue for model 17

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%



4.2.18 Score per residue for model 18

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 8% 5% 13% 74%



4.2.19 Score per residue for model 19

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



4.2.20 Score per residue for model 20

• Molecule 1: 1-ACYLGLYCEROL-3-PHOSPHATE O-ACYLTRANSFERASE ABHD5

Chain A: 5% 8% 13% 74%



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *LOWEST ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	refinement	
NMRDraw	structure solution	ANY
NMRPipe	structure solution	ANY
CcpNmr Analysis	structure solution	2.4
TALOS	structure solution	1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	124
Number of shifts mapped to atoms	124
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	46%

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	76	70	70	7±1
All	All	1520	1400	1400	143

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 49.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:25:TRP:CG	1:A:25:TRP:O	0.66	2.49	11	20
1:A:21:TRP:O	1:A:25:TRP:CD1	0.59	2.56	6	20
1:A:20:GLY:O	1:A:23:THR:HG23	0.56	2.00	2	20
1:A:21:TRP:CD1	1:A:22:LEU:HD22	0.53	2.38	15	20
1:A:26:LEU:HD13	1:A:26:LEU:N	0.53	2.19	17	12
1:A:18:GLY:HA2	1:A:22:LEU:HD23	0.48	1.85	1	19
1:A:21:TRP:HD1	1:A:22:LEU:HD22	0.45	1.71	15	20
1:A:25:TRP:H	1:A:26:LEU:HD13	0.42	1.75	20	11
1:A:26:LEU:HD23	1:A:26:LEU:N	0.41	2.31	8	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	10/39 (26%)	4±0 (40±2%)	5±0 (50±2%)	1±0 (10±0%)	1 10
All	All	200/780 (26%)	79 (40%)	101 (50%)	20 (10%)	1 10

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	20	GLY	20

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	7/28 (25%)	3±1 (37±8%)	4±1 (63±8%)	0 0
All	All	140/560 (25%)	52 (37%)	88 (63%)	0 0

All 5 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	21	TRP	20
1	A	22	LEU	20
1	A	25	TRP	20
1	A	26	LEU	19
1	A	19	SER	9

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 46% for the well-defined parts and 28% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chemical_shifts_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	124
Number of shifts mapped to atoms	124
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 46%, i.e. 58 atoms were assigned a chemical shift out of a possible 125. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	19/51 (37%)	19/22 (86%)	0/20 (0%)	0/9 (0%)
Sidechain	33/50 (66%)	33/34 (97%)	0/16 (0%)	0/0 (—%)
Aromatic	6/24 (25%)	6/12 (50%)	0/10 (0%)	0/2 (0%)
Overall	58/125 (46%)	58/68 (85%)	0/46 (0%)	0/11 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 28%, i.e. 124 atoms were assigned a chemical shift out of a possible 443. 0 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	41/198 (21%)	41/83 (49%)	0/78 (0%)	0/37 (0%)
Sidechain	74/202 (37%)	74/133 (56%)	0/67 (0%)	0/2 (0%)
Aromatic	9/43 (21%)	9/22 (41%)	0/17 (0%)	0/4 (0%)
Overall	124/443 (28%)	124/238 (52%)	0/162 (0%)	0/43 (0%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A: