



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2024 – 02:20 pm BST

PDB ID : 4AKL
Title : Structure of the Crimean-Congo Haemorrhagic Fever Virus Nucleocapsid Protein
Authors : Carter, S.D.; Walter, C.T.; Surtees, R.; Bergeron, E.; Ariza, A.; Albarino, C.G.; Nichol, S.T.; Hiscox, J.A.; Edwards, T.A.; Barr, J.N.
Deposited on : 2012-02-24
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	<div> <div>6%</div> <div> <div></div> <div>90%</div> <div>7%</div> <div>..</div> </div> </div>
1	B	482	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>.</div> </div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NUCLEOCAPSID.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	473	Total	C	N	O	S	0	16	0
			3830	2449	650	711	20			
1	B	473	Total	C	N	O	S	0	15	0
			3822	2446	648	708	20			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	111	ILE	THR	conflict	UNP Q70UR4
A	195	HIS	ARG	conflict	UNP Q70UR4
A	445	ASP	HIS	conflict	UNP Q70UR4
B	111	ILE	THR	conflict	UNP Q70UR4
B	195	HIS	ARG	conflict	UNP Q70UR4
B	445	ASP	HIS	conflict	UNP Q70UR4

- Molecule 2 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	6	4		
2	B	1	Total	C	O	0	0
			10	6	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	151	Total	O	0	0
			151	151		
3	B	162	Total	O	0	0
			162	162		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NUCLEOCAPSID



• Molecule 1: NUCLEOCAPSID



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	150.38Å 72.06Å 101.23Å 90.00° 110.70° 90.00°	Depositor
Resolution (Å)	94.70 – 2.10 47.90 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.2 (94.70-2.10) 95.2 (47.90-2.10)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.184 , 0.234 0.184 , 0.234	Depositor DCC
R_{free} test set	2857 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtriage
Anisotropy	0.283	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7985	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	2/3959 (0.1%)	0.76	4/5335 (0.1%)
1	B	1.06	2/3948 (0.1%)	0.81	4/5321 (0.1%)
All	All	1.02	4/7907 (0.1%)	0.79	8/10656 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	74	CYS	CB-SG	-8.94	1.67	1.82
1	B	121	GLU	CG-CD	7.29	1.62	1.51
1	B	74	CYS	CB-SG	-6.86	1.70	1.82
1	A	404	ALA	CA-CB	5.10	1.63	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	372	ARG	NE-CZ-NH1	10.87	125.74	120.30
1	B	372	ARG	NE-CZ-NH1	10.71	125.66	120.30
1	A	372	ARG	NE-CZ-NH2	-7.21	116.70	120.30
1	B	134	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	B	372	ARG	NE-CZ-NH2	-6.50	117.05	120.30
1	A	428	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	B	401	ASP	CB-CG-OD1	5.36	123.12	118.30
1	A	384	ARG	NE-CZ-NH2	-5.07	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3830	0	3878	25	0
1	B	3822	0	3874	18	0
2	A	10	0	14	1	0
2	B	10	0	14	0	0
3	A	151	0	0	0	0
3	B	162	0	0	2	0
All	All	7985	0	7780	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:LEU:HD12	1:A:246:VAL:HG21	1.70	0.74
1:B:446:MET:HE1	3:B:2151:HOH:O	2.01	0.60
1:A:342[B]:LYS:NZ	1:B:369:GLN:HE22	1.99	0.60
1:B:330:PHE:O	1:B:334:LEU:HD13	2.02	0.60
1:A:475:ASN:OD1	1:A:477:THR:HG22	2.06	0.55
1:A:365:ASP:O	1:A:372:ARG:NH2	2.39	0.55
1:B:170:LEU:HD11	1:B:481:ILE:HD11	1.91	0.53
1:A:228:ILE:HD11	1:A:286:LYS:HE3	1.89	0.53
1:A:68:VAL:HG21	1:A:338:PRO:HB3	1.92	0.52
1:A:453[A]:HIS:CE1	1:A:457:GLN:NE2	2.78	0.52
1:A:202:ARG:O	1:A:206:LYS:HD2	2.10	0.51
1:A:433:LEU:O	1:A:437[A]:GLN:HG3	2.11	0.50
1:A:342[B]:LYS:HE3	1:A:342[B]:LYS:HA	1.95	0.49
1:B:44:ASP:OD1	1:B:432:LYS:HE3	2.13	0.49
1:B:63:TYR:CE1	1:B:437[A]:GLN:HG2	2.48	0.49
1:B:261:LEU:HD13	1:B:273:ALA:HB1	1.95	0.49
1:A:437[B]:GLN:NE2	1:A:446:MET:SD	2.86	0.47
1:B:71:THR:HG22	1:B:413:ILE:HG13	1.95	0.47
1:A:63:TYR:CE1	1:A:437[A]:GLN:HG2	2.49	0.47
1:A:248:ASP:OD1	1:A:248:ASP:N	2.44	0.47
1:A:202:ARG:HB2	1:A:238:LEU:HD21	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:ILE:HD11	1:A:290:LEU:HD21	1.98	0.46
1:A:342[B]:LYS:HA	1:A:342[B]:LYS:CE	2.46	0.45
1:B:155:GLU:HG2	1:B:482:ILE:HD11	1.99	0.45
1:B:247:PHE:CG	1:B:288:GLN:HG2	2.53	0.44
1:B:228:ILE:CG2	1:B:233:THR:HG23	2.48	0.44
1:A:117:VAL:HG11	2:A:1483:PGE:H42	2.00	0.43
1:A:372:ARG:HH11	1:A:372:ARG:CG	2.30	0.43
1:A:228:ILE:HD11	1:A:290:LEU:HD11	2.00	0.43
1:A:228:ILE:CD1	1:A:290:LEU:HD21	2.49	0.42
1:B:456:HIS:O	1:B:460:VAL:HG13	2.19	0.42
1:A:251[B]:LYS:HG3	1:A:284:VAL:HG11	2.01	0.42
1:A:342[B]:LYS:HZ2	1:B:369:GLN:HE22	1.67	0.42
1:A:372:ARG:HH11	1:A:372:ARG:HG2	1.85	0.41
1:B:149:SER:O	1:B:473[A]:LYS:NZ	2.54	0.41
1:B:174:ILE:HG12	1:B:477:THR:HG23	2.02	0.41
1:B:475:ASN:OD1	1:B:477:THR:HB	2.20	0.41
1:B:358:LYS:HD2	3:B:2112:HOH:O	2.20	0.40
1:A:460:VAL:HG12	1:A:476:ALA:HB2	2.04	0.40
1:B:457:GLN:HB3	1:B:462:LYS:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/482 (101%)	479 (99%)	6 (1%)	0	100	100
1	B	484/482 (100%)	473 (98%)	11 (2%)	0	100	100
All	All	969/964 (100%)	952 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	418/408 (102%)	405 (97%)	13 (3%)	40	43
1	B	417/408 (102%)	406 (97%)	11 (3%)	46	50
All	All	835/816 (102%)	811 (97%)	24 (3%)	42	46

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LYS
1	A	102	THR
1	A	164[A]	MET
1	A	164[B]	MET
1	A	206	LYS
1	A	238	LEU
1	A	241	THR
1	A	248	ASP
1	A	339	ARG
1	A	372	ARG
1	A	423	ASN
1	A	473	LYS
1	A	477	THR
1	B	19	GLU
1	B	55	ASP
1	B	181	LEU
1	B	223	SER
1	B	225	ARG
1	B	241	THR
1	B	248	ASP
1	B	265	LYS
1	B	270	LYS
1	B	372	ARG
1	B	457	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	288	GLN
1	A	419	ASN
1	A	457	GLN
1	A	463	GLN
1	B	182	ASN
1	B	369	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PGE	A	1483	-	9,9,9	0.78	0	8,8,8	1.03	1 (12%)
2	PGE	B	1483	-	9,9,9	0.86	0	8,8,8	1.12	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PGE	A	1483	-	-	3/7/7/7	-
2	PGE	B	1483	-	-	4/7/7/7	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1483	PGE	O4-C6-C5	-2.82	95.43	111.81
2	A	1483	PGE	O4-C6-C5	-2.47	97.49	111.81

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1483	PGE	O2-C3-C4-O3
2	A	1483	PGE	O2-C3-C4-O3
2	B	1483	PGE	O1-C1-C2-O2
2	A	1483	PGE	C1-C2-O2-C3
2	B	1483	PGE	C1-C2-O2-C3
2	B	1483	PGE	C6-C5-O3-C4
2	A	1483	PGE	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1483	PGE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	473/482 (98%)	0.20	31 (6%) 18 23	25, 45, 87, 145	0
1	B	473/482 (98%)	0.01	11 (2%) 60 65	24, 42, 63, 84	0
All	All	946/964 (98%)	0.11	42 (4%) 34 40	24, 44, 74, 145	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	225	ARG	6.3
1	A	225	ARG	6.2
1	A	227	GLY	6.0
1	A	226	SER	5.8
1	A	192	PRO	5.3
1	A	228	ILE	5.1
1	A	290	LEU	4.7
1	B	226	SER	4.6
1	A	193	VAL	4.4
1	B	224	GLY	4.2
1	A	224	GLY	3.9
1	A	289	GLU	3.9
1	A	195	HIS	3.8
1	A	230	LEU	3.8
1	B	192	PRO	3.7
1	A	287	ALA	3.6
1	B	228	ILE	3.4
1	B	193	VAL	3.4
1	B	223	SER	3.3
1	A	223	SER	3.3
1	A	247	PHE	3.2
1	A	246	VAL	3.1
1	A	233	THR	2.8
1	A	179	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	244	LYS	2.7
1	A	285	ALA	2.7
1	A	241	THR	2.6
1	A	196	GLU	2.6
1	B	241	THR	2.6
1	A	291	TYR	2.5
1	A	282[A]	LYS	2.5
1	B	219	ASP	2.2
1	A	229	ALA	2.2
1	A	242	GLU	2.2
1	A	237	LYS	2.1
1	A	198	VAL	2.1
1	A	243	GLY	2.1
1	B	282[A]	LYS	2.1
1	A	281	LEU	2.1
1	A	234	GLY	2.1
1	A	251[A]	LYS	2.1
1	B	245	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PGE	A	1483	10/10	0.75	0.26	58,59,60,61	0
2	PGE	B	1483	10/10	0.86	0.19	50,54,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.