



# Full wwPDB NMR Structure Validation Report ⓘ

May 7, 2024 – 03:59 pm BST

PDB ID : 2BGF  
Title : NMR structure of Lys48-linked di-ubiquitin using chemical shift perturbation data together with RDCs and 15N-relaxation data  
Authors : Van Dijk, A.D.J.; Fushman, D.; Bonvin, A.M.J.J.  
Deposited on : 2004-12-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2

## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

There are no overall percentile quality scores available for this entry.

The sequence quality summary graphics cannot be shown.

## 2 Ensemble composition and analysis

This entry contains 10 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:72, B:1-B:72 (144)	0.90	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 5, 6, 7, 8, 9, 10
Single-model clusters	4

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2460 atoms, of which 1257 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DI-UBIQUITIN.


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	76	1231	378	630	105	117	1	0
1	B	76	1229	378	627	105	118	1	0

## 4 Residue-property plots


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DI-UBIQUITIN



- Molecule 1: DI-UBIQUITIN




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: DI-UBIQUITIN




- Molecule 1: DI-UBIQUITIN




### 4.2.2 Score per residue for model 2

- Molecule 1: DI-UBIQUITIN




- Molecule 1: DI-UBIQUITIN




### 4.2.3 Score per residue for model 3

- Molecule 1: DI-UBIQUITIN




- Molecule 1: DI-UBIQUITIN




### 4.2.4 Score per residue for model 4

- Molecule 1: DI-UBIQUITIN




- Molecule 1: DI-UBIQUITIN




#### 4.2.5 Score per residue for model 5

- Molecule 1: DI-UBIQUITIN




- Molecule 1: DI-UBIQUITIN




#### 4.2.6 Score per residue for model 6

- Molecule 1: DI-UBIQUITIN




- Molecule 1: DI-UBIQUITIN




#### 4.2.7 Score per residue for model 7 (medoid)

- Molecule 1: DI-UBIQUITIN




- Molecule 1: DI-UBIQUITIN




#### 4.2.8 Score per residue for model 8

- Molecule 1: DI-UBIQUITIN

Chain A:  95% 5%




- Molecule 1: DI-UBIQUITIN

Chain B:  95% 5%




#### 4.2.9 Score per residue for model 9

- Molecule 1: DI-UBIQUITIN

Chain A:  95% 5%




- Molecule 1: DI-UBIQUITIN

Chain B:  95% 5%




#### 4.2.10 Score per residue for model 10

- Molecule 1: DI-UBIQUITIN

Chain A:  95% 5%



- Molecule 1: DI-UBIQUITIN

Chain B:  95% 5%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *HADDOCK*.

Of the 200 calculated structures, 10 were deposited, based on the following criterion: *LOWEST ENERGY STRUCTURES OF LOWEST ENERGY CLUSTER*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
HADDOCK	refinement	
HADDOCK	structure solution	

No chemical shift data was provided.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

MolProbity failed to run properly - this section will have to be empty.

### 6.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section will have to be empty.

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section will have to be empty.

#### 6.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section will have to be empty.

#### 6.3.3 RNA [i](#)

MolProbity failed to run properly - this section will have to be empty.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

MolProbity failed to run properly - this section will have to be empty.

### 6.5 Carbohydrates [i](#)

MolProbity failed to run properly - this section will have to be empty.

### 6.6 Ligand geometry [i](#)

MolProbity failed to run properly - this section will have to be empty.

### 6.7 Other polymers [i](#)

MolProbity failed to run properly - this section will have to be empty.

## 6.8 Polymer linkage issues

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided