



# wwPDB X-ray Structure Validation Summary Report

Feb 27, 2023 – 06:30 pm GMT

PDB ID : 8BLT  
Title : Structure of Lactobacillus salivarius (Ls) bile salt hydrolase(BSH) in complex with taurocholate (TCA)  
Authors : Karlov, D.S.; Long, S.L.; Zeng, X.; Xu, F.; Lal, K.; Cao, L.; Hayoun, K.; Lin, J.; Joyce, S.A.; Tikhonova, I.G.  
Deposited on : 2022-11-10  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.32.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	 91% 6% ..
1	B	325	 93% 6% .
1	C	325	 93% 6% ..
1	D	325	 88% 9% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	E	325	 2% 91% 7% .
1	F	325	 2% 89% 9% .
1	G	325	 7% 89% 8% ..
1	H	325	 4% 89% 10% .

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 21507 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bile salt hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	321	Total 2546	C 1622	N 417	O 501	S 6	0	0	0
1	B	321	Total 2550	C 1625	N 417	O 502	S 6	0	1	0
1	C	321	Total 2551	C 1623	N 420	O 502	S 6	0	0	0
1	D	319	Total 2508	C 1598	N 411	O 493	S 6	0	0	0
1	E	318	Total 2477	C 1579	N 405	O 487	S 6	0	0	0
1	F	318	Total 2519	C 1603	N 410	O 500	S 6	0	0	0
1	G	318	Total 2506	C 1598	N 412	O 490	S 6	0	0	0
1	H	319	Total 2532	C 1612	N 416	O 498	S 6	0	0	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	325	LEU	-	expression tag	UNP J7H3P9
A	326	GLU	-	expression tag	UNP J7H3P9
B	325	LEU	-	expression tag	UNP J7H3P9
B	326	GLU	-	expression tag	UNP J7H3P9
C	325	LEU	-	expression tag	UNP J7H3P9
C	326	GLU	-	expression tag	UNP J7H3P9
D	325	LEU	-	expression tag	UNP J7H3P9
D	326	GLU	-	expression tag	UNP J7H3P9
E	325	LEU	-	expression tag	UNP J7H3P9
E	326	GLU	-	expression tag	UNP J7H3P9
F	325	LEU	-	expression tag	UNP J7H3P9
F	326	GLU	-	expression tag	UNP J7H3P9
G	325	LEU	-	expression tag	UNP J7H3P9

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	326	GLU	-	expression tag	UNP J7H3P9
H	325	LEU	-	expression tag	UNP J7H3P9
H	326	GLU	-	expression tag	UNP J7H3P9

- Molecule 2 is TAUROCHOLIC ACID (three-letter code: TCH) (formula:  $C_{26}H_{45}NO_7S$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
2	B	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
2	C	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
2	D	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
2	E	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
2	F	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
2	G	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
2	H	1	Total	C	N	O	S	0	0
			35	26	1	7	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	155	Total 161	O 161	0	6
3	B	195	Total 198	O 198	0	3
3	C	179	Total 181	O 181	0	2
3	D	119	Total 119	O 119	0	0
3	E	98	Total 98	O 98	0	0
3	F	98	Total 100	O 100	0	2
3	G	92	Total 93	O 93	0	1
3	H	88	Total 88	O 88	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bile salt hydrolase




- Molecule 1: Bile salt hydrolase



- Molecule 1: Bile salt hydrolase




- Molecule 1: Bile salt hydrolase





- Molecule 1: Bile salt hydrolase









• Molecule 1: Bile salt hydrolase



• Molecule 1: Bile salt hydrolase



• Molecule 1: Bile salt hydrolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.76Å 94.61Å 168.22Å 90.00° 90.26° 90.00°	Depositor
Resolution (Å)	25.75 – 2.10 25.74 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.6 (25.75-2.10) 96.7 (25.74-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.05 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.180 , 0.213 0.187 , 0.218	Depositor DCC
$R_{free}$ test set	7772 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.9	Xtrriage
Anisotropy	0.029	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.033 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	21507	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TCH, OCS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.75	0/2590	0.89	1/3513 (0.0%)
1	B	0.77	0/2594	0.88	0/3520
1	C	0.78	0/2595	0.91	0/3520
1	D	0.73	0/2552	0.89	1/3468 (0.0%)
1	E	0.73	0/2520	0.87	0/3426
1	F	0.73	0/2563	0.86	0/3481
1	G	0.76	1/2550 (0.0%)	0.87	0/3463
1	H	0.73	0/2576	0.87	0/3494
All	All	0.75	1/20540 (0.0%)	0.88	2/27885 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	121	GLU	CD-OE1	-5.65	1.19	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	26	GLU	CB-CA-C	-6.57	97.26	110.40
1	D	26	GLU	CB-CA-C	-5.39	99.62	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2546	0	2463	15	0
1	B	2550	0	2466	6	0
1	C	2551	0	2471	12	0
1	D	2508	0	2404	19	0
1	E	2477	0	2362	8	0
1	F	2519	0	2421	15	0
1	G	2506	0	2417	14	0
1	H	2532	0	2450	20	0
2	A	35	0	44	2	0
2	B	35	0	44	1	0
2	C	35	0	44	3	0
2	D	35	0	44	5	0
2	E	35	0	44	0	0
2	F	35	0	44	5	0
2	G	35	0	44	6	0
2	H	35	0	44	7	0
3	A	161	0	0	2	0
3	B	198	0	0	0	0
3	C	181	0	0	1	0
3	D	119	0	0	3	0
3	E	98	0	0	1	0
3	F	100	0	0	3	0
3	G	93	0	0	3	0
3	H	88	0	0	0	0
All	All	21507	0	19806	123	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 123 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:235:SER:HB2	1:F:247:GLN:HE21	1.41	0.84
2:F:401:TCH:H26	3:F:503:HOH:O	1.89	0.71
1:C:79:ASN:ND2	2:C:401:TCH:H26A	2.07	0.70
2:G:401:TCH:C26	3:G:509:HOH:O	2.40	0.68
2:F:401:TCH:C26	3:F:503:HOH:O	2.41	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/325 (98%)	306 (96%)	11 (4%)	0	100	100
1	B	318/325 (98%)	312 (98%)	6 (2%)	0	100	100
1	C	317/325 (98%)	309 (98%)	8 (2%)	0	100	100
1	D	315/325 (97%)	301 (96%)	14 (4%)	0	100	100
1	E	314/325 (97%)	303 (96%)	11 (4%)	0	100	100
1	F	314/325 (97%)	306 (98%)	8 (2%)	0	100	100
1	G	314/325 (97%)	303 (96%)	11 (4%)	0	100	100
1	H	315/325 (97%)	306 (97%)	9 (3%)	0	100	100
All	All	2524/2600 (97%)	2446 (97%)	78 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/287 (97%)	271 (98%)	7 (2%)	47	52
1	B	279/287 (97%)	271 (97%)	8 (3%)	42	46
1	C	280/287 (98%)	274 (98%)	6 (2%)	53	59
1	D	271/287 (94%)	265 (98%)	6 (2%)	52	57
1	E	265/287 (92%)	257 (97%)	8 (3%)	41	44
1	F	276/287 (96%)	270 (98%)	6 (2%)	52	57

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	272/287 (95%)	267 (98%)	5 (2%)	59	65
1	H	277/287 (96%)	272 (98%)	5 (2%)	59	65
All	All	2198/2296 (96%)	2147 (98%)	51 (2%)	50	55

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	24	TYR
1	F	16	ARG
1	H	140	HIS
1	E	94	LYS
1	E	140	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	187	ASN
1	F	247	GLN
1	H	294	GLN
1	G	247	GLN
1	C	133	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	OCS	C	2	1	7,8,9	0.79	0	6,11,13	2.02	1 (16%)
1	OCS	G	2	1	7,8,9	1.27	1 (14%)	6,11,13	1.35	0
1	OCS	F	2	1	7,8,9	1.17	1 (14%)	6,11,13	2.96	2 (33%)
1	OCS	E	2	1	7,8,9	1.25	1 (14%)	6,11,13	5.60	4 (66%)
1	OCS	H	2	1	7,8,9	1.09	1 (14%)	6,11,13	2.49	3 (50%)
1	OCS	D	2	1	7,8,9	0.91	0	6,11,13	2.06	3 (50%)
1	OCS	B	2	1	7,8,9	1.21	1 (14%)	6,11,13	1.33	1 (16%)
1	OCS	A	2	1	7,8,9	1.48	1 (14%)	6,11,13	2.21	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	C	2	1	-	0/4/7/9	-
1	OCS	G	2	1	-	0/4/7/9	-
1	OCS	F	2	1	-	0/4/7/9	-
1	OCS	E	2	1	-	1/4/7/9	-
1	OCS	H	2	1	-	0/4/7/9	-
1	OCS	D	2	1	-	0/4/7/9	-
1	OCS	B	2	1	-	0/4/7/9	-
1	OCS	A	2	1	-	1/4/7/9	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	OCS	OD3-SG	3.44	1.55	1.45
1	G	2	OCS	OD3-SG	2.84	1.53	1.45
1	E	2	OCS	OD3-SG	2.81	1.53	1.45
1	F	2	OCS	OD1-SG	2.57	1.52	1.45
1	B	2	OCS	OD3-SG	2.29	1.51	1.45

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	2	OCS	OD3-SG-CB	-12.66	91.89	106.94
1	F	2	OCS	OD1-SG-CB	-6.79	98.87	106.94
1	H	2	OCS	OD2-SG-OD3	4.59	122.49	111.27
1	C	2	OCS	OD3-SG-CB	-4.54	101.55	106.94

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	OCS	OD3-SG-CB	-3.68	102.57	106.94

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	2	OCS	N-CA-CB-SG
1	E	2	OCS	CA-CB-SG-OD1

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	G	2	OCS	1	0
1	A	2	OCS	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TCH	A	401	-	38,38,38	0.48	0	59,60,60	0.91	2 (3%)
2	TCH	B	401	-	38,38,38	0.51	0	59,60,60	1.05	2 (3%)
2	TCH	D	401	-	38,38,38	0.56	0	59,60,60	1.00	2 (3%)
2	TCH	F	401	-	38,38,38	0.58	0	59,60,60	0.93	2 (3%)
2	TCH	E	401	-	38,38,38	0.44	0	59,60,60	0.83	2 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TCH	H	401	-	38,38,38	0.71	0	59,60,60	1.34	8 (13%)
2	TCH	G	401	-	38,38,38	0.45	0	59,60,60	1.11	4 (6%)
2	TCH	C	401	-	38,38,38	0.70	0	59,60,60	1.27	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TCH	A	401	-	-	6/16/81/81	0/4/4/4
2	TCH	B	401	-	-	6/16/81/81	0/4/4/4
2	TCH	D	401	-	-	6/16/81/81	0/4/4/4
2	TCH	F	401	-	-	2/16/81/81	0/4/4/4
2	TCH	E	401	-	-	3/16/81/81	0/4/4/4
2	TCH	H	401	-	-	11/16/81/81	0/4/4/4
2	TCH	G	401	-	-	6/16/81/81	0/4/4/4
2	TCH	C	401	-	-	4/16/81/81	0/4/4/4

There are no bond length outliers.

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	TCH	O1S-S26-C26	-4.76	101.19	106.92
2	H	401	TCH	C22-C23-C24	4.02	122.02	113.04
2	G	401	TCH	C9-C8-C7	3.74	116.35	111.88
2	C	401	TCH	O3S-S26-C26	3.29	111.09	105.77
2	E	401	TCH	O2S-S26-C26	-3.07	103.22	106.92

There are no chirality outliers.

5 of 44 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	TCH	C25-C26-S26-O1S
2	A	401	TCH	C25-C26-S26-O3S
2	B	401	TCH	C20-C22-C23-C24
2	B	401	TCH	C23-C24-N24-C25
2	B	401	TCH	C25-C26-S26-O1S




There are no ring outliers.




7 monomers are involved in 29 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	TCH	2	0
2	B	401	TCH	1	0
2	D	401	TCH	5	0
2	F	401	TCH	5	0
2	H	401	TCH	7	0
2	G	401	TCH	6	0
2	C	401	TCH	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	320/325 (98%)	-0.24	2 (0%) 89 91	16, 26, 48, 67	0
1	B	320/325 (98%)	-0.38	4 (1%) 77 80	14, 21, 40, 63	0
1	C	320/325 (98%)	-0.38	3 (0%) 84 86	15, 21, 38, 66	0
1	D	318/325 (97%)	-0.09	6 (1%) 66 71	16, 30, 50, 71	0
1	E	317/325 (97%)	0.06	7 (2%) 62 66	17, 36, 58, 68	0
1	F	317/325 (97%)	-0.06	5 (1%) 72 75	17, 31, 52, 67	0
1	G	317/325 (97%)	0.21	23 (7%) 15 19	18, 36, 55, 68	0
1	H	318/325 (97%)	0.09	14 (4%) 34 40	18, 34, 53, 92	0
All	All	2547/2600 (97%)	-0.10	64 (2%) 57 62	14, 29, 53, 92	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	305	ASP	4.4
1	H	324	ASN	4.3
1	G	103	ILE	3.9
1	E	103	ILE	3.8
1	F	92	ASN	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	OCS	G	2	9/10	0.96	0.10	23,28,43,44	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	OCS	E	2	9/10	0.97	0.08	23,28,40,46	0
1	OCS	C	2	9/10	0.98	0.11	17,20,30,31	0
1	OCS	D	2	9/10	0.98	0.10	20,26,35,38	0
1	OCS	A	2	9/10	0.98	0.08	19,22,27,28	0
1	OCS	F	2	9/10	0.98	0.10	23,26,35,43	0
1	OCS	B	2	9/10	0.98	0.09	15,23,30,34	0
1	OCS	H	2	9/10	0.98	0.09	21,26,34,36	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	TCH	G	401	35/35	0.83	0.17	44,56,73,79	0
2	TCH	H	401	35/35	0.87	0.19	40,65,97,100	0
2	TCH	F	401	35/35	0.88	0.14	34,46,69,69	0
2	TCH	C	401	35/35	0.89	0.14	25,31,65,69	0
2	TCH	A	401	35/35	0.90	0.15	34,47,54,55	0
2	TCH	E	401	35/35	0.91	0.14	38,48,66,69	0
2	TCH	D	401	35/35	0.91	0.12	36,40,58,61	0
2	TCH	B	401	35/35	0.92	0.10	25,27,59,61	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


**Electron density around TCH G 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TCH H 401:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around TCH A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around TCH E 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around TCH D 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TCH B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.