Full wwPDB X-ray Structure Validation Report (i)

May 27, 2020 – 03:29 am BST

PDB ID	:	5C86
Title	:	Novel fungal alcohol oxidase with catalytic diversity among the AA5 family, apo form
Authors	:	Urresti, S.; Yin, D.T.; LaFond, M.; Derikvand, F.; Berrin, G.J.; Henrissat, B.; Walton, P.H.; Brumer, H.; Davies, G.J.
Deposited on	:	2015-06-25
Resolution	:	1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	89% 9% •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kelch domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	482	3767	2382	641	716	28	0	19	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	483	HIS	-	expression tag	UNP E3QHV8
A	484	HIS	-	expression tag	UNP E3QHV8
A	485	HIS	-	expression tag	UNP E3QHV8
A	486	HIS	-	expression tag	UNP E3QHV8
A	487	HIS	-	expression tag	UNP E3QHV8
A	488	HIS	-	expression tag	UNP E3QHV8

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	403	403	403	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kelch domain-containing protein

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α, β, γ	123.71Å 123.71Å 52.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.53 – 1.51 40.49 – 1.51	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.53-1.51) 99.9 (40.49-1.51)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.03	Depositor
$< I/\sigma(I) >$ [1]	3.20 (at 1.51Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.150 , 0.183 0.161 , 0.190	Depositor DCC
R_{free} test set	3581 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å^2)	15.6	Xtriage
Anisotropy	0.774	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.31 , 38.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	4170	wwPDB-VP
Average B, all atoms (Å^2)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	1.12	5/3871 (0.1%)	1.12	18/5272 (0.3%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	96	SER	CA-CB	-6.72	1.42	1.52
1	A	411	TYR	CE1-CZ	-5.69	1.31	1.38
1	A	184	TRP	CG-CD1	5.27	1.44	1.36
1	A	6	TRP	CE3-CZ3	5.15	1.47	1.38
1	A	104	GLU	CD-OE2	-5.03	1.20	1.25

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	443	LYS	CD-CE-NZ	10.21	135.20	111.70
1	A	171	ARG	NE-CZ-NH1	8.11	124.35	120.30
1	A	79	ASP	CB-CG-OD1	7.47	125.03	118.30
1	A	118	ARG	NE-CZ-NH1	7.44	124.02	120.30
1	A	358[A]	CYS	CA-CB-SG	-7.21	101.03	114.00
1	A	358[B]	CYS	CA-CB-SG	-7.21	101.03	114.00
1	A	316	ASP	CB-CG-OD2	-7.18	111.83	118.30
1	A	372	TYR	CB-CG-CD1	7.01	125.20	121.00
1	A	171	ARG	NE-CZ-NH2	-6.36	117.12	120.30
1	A	70	ASP	CB-CG-OD2	6.00	123.70	118.30
1	A	407	ASP	CB-CG-OD1	-5.89	113.00	118.30
1	A	176	ASP	CB-CG-OD1	5.68	123.41	118.30
1	A	438	ASP	CB-CG-OD1	5.54	123.29	118.30
1	A	79	ASP	CB-CG-OD2	-5.49	113.36	118.30
1	A	428	ASP	CB-CG-OD1	5.25	123.02	118.30
1	A	257	LEU	CB-CG-CD2	5.19	119.82	111.00
1	A	38[A]	ARG	NE-CZ-NH1	-5.08	117.76	120.30
1	A	38[B]	ARG	NE-CZ-NH1	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3767	0	3645	21	0
2	A	403	0	0	2	1
All	All	4170	0	3645	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9[A]:MET:HE2	1:A:364:ASP:OD2	1.82	0.79
1:A:9[A]:MET:CE	1:A:364:ASP:OD2	2.29	0.79
1:A:173[A]:MET:HE2	2:A:815:HOH:O	1.93	0.69
1:A:153:ALA:HB3	1:A:155[B]:THR:HG22	1.86	0.58
1:A:9[A]:MET:HE1	1:A:364:ASP:OD2	2.09	0.53
1:A:256[A]:ILE:HG12	1:A:271:LEU:HD11	1.92	0.51
1:A:38[A]:ARG:HB3	1:A:355:GLY:HA2	1.93	0.51
1:A:34:GLY:HA3	1:A:44:ASN:HB3	1.93	0.50
1:A:9[A]:MET:HE3	1:A:365:GLY:O	2.12	0.50
1:A:328:HIS:HD2	1:A:330:VAL:O	1.95	0.49
1:A:403:ILE:HG12	1:A:463[A]:LEU:HD21	1.94	0.49
1:A:287:PRO:HA	1:A:416:ILE:HD11	1.95	0.48
1:A:227[B]:VAL:HG11	1:A:285:VAL:HB	1.97	0.46
1:A:396:ARG:HD3	1:A:483:HIS:CE1	2.50	0.45
1:A:155[B]:THR:HG23	1:A:155[B]:THR:O	2.17	0.45
1:A:390:LEU:HD21	1:A:463[A]:LEU:HD23	1.99	0.44
1:A:9[A]:MET:HE2	1:A:364:ASP:HB2	1.99	0.44
1:A:73:CYS:N	1:A:74:PRO:HD3	2.34	0.43
1:A:328:HIS:HE1	2:A:650:HOH:O	2.01	0.43
1:A:247:GLY:O	1:A:299:LYS:HE3	2.19	0.41
1:A:19:VAL:HG12	1:A:30:VAL:HG12	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:567:HOH:O	2:A:610:HOH:O[4_664]	2.15	0.05

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	499/488 (102%)	479 (96%)	20 (4%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	406/393 (103%)	405 (100%)	1 (0%)	93 86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	372	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	ASN
1	A	58	ASN
1	A	115	GLN
1	A	269	GLN
1	A	328	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	482/488 (98%)	-0.53	2 (0%) 92 94	10, 17, 27, 51	10 (2%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	483	HIS	2.4
1	A	357	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.