



## wwPDB EM Validation Summary Report i

Feb 11, 2024 – 06:20 AM EST

PDB ID : 3C92  
EMDB ID : EMD-1740  
Title : Thermoplasma acidophilum 20S proteasome with a closed gate  
Authors : Rabl, J.; Smith, D.M.; Yu, Y.; Chang, S.C.; Goldberg, A.L.; Cheng, Y.  
Deposited on : 2008-02-14  
Resolution : 6.80 Å(reported)  
Based on initial models : 3C91, 1PMA

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70  
MolProbit : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*


The reported resolution of this entry is 6.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.



*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain			
1	P	233	35%	48%	42%	6% ..
1	Q	233	29%	47%	42%	6% ..
1	R	233	31%	47%	42%	6% ..
1	S	233	29%	46%	43%	6% ..
1	T	233	29%	47%	42%	6% ..
1	U	233	33%	48%	41%	6% ..
2	1	203	25%	59%	34%	7%
2	2	203	23%	55%	37%	8%
2	H	203	23%	53%	38%	9%
2	I	203	24%	61%	31%	8%
2	J	203	24%	60%	32%	8%
2	K	203	25%	54%	38%	8%
2	L	203	24%	53%	38%	8%
2	M	203	22%	58%	35%	7%
2	N	203	24%	54%	39%	7%
2	V	203	27%	53%	38%	9%
2	W	203	27%	60%	32%	8%
2	X	203	26%	60%	33%	7%
2	Y	203	25%	52%	40%	8%
2	Z	203	25%	54%	38%	8%

## 2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 46228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	B	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	C	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	D	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	E	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	F	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	G	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	O	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	P	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	Q	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	R	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	S	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	T	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		
1	U	224	Total	C	N	O	S	0	0
			1744	1107	296	338	3		

- Molecule 2 is a protein called Proteasome subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	H	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
2	I	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	J	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	K	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	L	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	M	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	N	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	V	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	W	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	X	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	Y	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	Z	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	1	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0
2	2	203	Total	C	N	O	S		
			1558	985	264	298	11	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha






- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha






- Molecule 1: Proteasome subunit alpha

Chain P: 35% • 48% • 42% • 6% •




- Molecule 1: Proteasome subunit alpha


Chain Q: 29% (Red), 47% (Green), 42% (Yellow), 6% (Black)




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha










- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta






- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



## 4 Experimental information i

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, D7	Depositor
Number of particles used	41546	Depositor
Resolution determination method	Not provided	
CTF correction method	PHASE AND AMPLITUDE	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	20	Depositor
Minimum defocus (nm)	-1500.00	Depositor
Maximum defocus (nm)	-3500.00	Depositor
Magnification	51159	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	8.466	Depositor
Minimum map value	-4.027	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	2.5	Depositor
Map size (Å)	197.028, 197.028, 197.028	wwPDB
Map dimensions	144, 144, 144	wwPDB
Map angles (°)	90, 90, 90	wwPDB
Pixel spacing (Å)	1.36825, 1.36825, 1.36825	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.55	0/1767	0.81	6/2380 (0.3%)
1	B	0.55	0/1767	0.81	6/2380 (0.3%)
1	C	0.55	0/1767	0.81	6/2380 (0.3%)
1	D	0.55	0/1767	0.81	6/2380 (0.3%)
1	E	0.55	0/1767	0.81	6/2380 (0.3%)
1	F	0.56	0/1767	0.81	6/2380 (0.3%)
1	G	0.56	0/1767	0.81	6/2380 (0.3%)
1	O	0.55	0/1767	0.81	6/2380 (0.3%)
1	P	0.56	0/1767	0.81	6/2380 (0.3%)
1	Q	0.56	0/1767	0.81	6/2380 (0.3%)
1	R	0.55	0/1767	0.81	6/2380 (0.3%)
1	S	0.55	0/1767	0.81	6/2380 (0.3%)
1	T	0.56	0/1767	0.82	6/2380 (0.3%)
1	U	0.56	0/1767	0.81	6/2380 (0.3%)
2	1	0.55	0/1577	0.75	1/2129 (0.0%)
2	2	0.54	0/1577	0.75	0/2129
2	H	0.57	0/1577	0.76	0/2129
2	I	0.57	0/1577	0.76	0/2129
2	J	0.54	0/1577	0.75	0/2129
2	K	0.55	0/1577	0.77	2/2129 (0.1%)
2	L	0.57	0/1577	0.77	0/2129
2	M	0.55	0/1577	0.75	1/2129 (0.0%)
2	N	0.54	0/1577	0.75	0/2129
2	V	0.57	0/1577	0.76	0/2129
2	W	0.57	0/1577	0.76	0/2129
2	X	0.54	0/1577	0.75	0/2129
2	Y	0.55	0/1577	0.77	2/2129 (0.1%)
2	Z	0.57	0/1577	0.77	0/2129
All	All	0.56	0/46816	0.79	90/63126 (0.1%)

There are no bond length outliers.

The worst 5 of 90 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	R	12	ILE	CB-CA-C	-10.42	90.76	111.60
1	T	12	ILE	CB-CA-C	-10.41	90.79	111.60
1	C	12	ILE	CB-CA-C	-10.40	90.79	111.60
1	G	12	ILE	CB-CA-C	-10.40	90.79	111.60
1	U	12	ILE	CB-CA-C	-10.40	90.80	111.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1744	0	1781	93	0
1	B	1744	0	1781	90	0
1	C	1744	0	1781	91	0
1	D	1744	0	1781	95	0
1	E	1744	0	1781	100	0
1	F	1744	0	1781	93	0
1	G	1744	0	1781	95	0
1	O	1744	0	1781	91	0
1	P	1744	0	1781	91	0
1	Q	1744	0	1781	93	0
1	R	1744	0	1781	97	0
1	S	1744	0	1781	100	0
1	T	1744	0	1781	95	0
1	U	1744	0	1781	94	0
2	1	1558	0	1609	80	0
2	2	1558	0	1609	74	0
2	H	1558	0	1609	81	0
2	I	1558	0	1609	62	0
2	J	1558	0	1609	65	0
2	K	1558	0	1609	71	0
2	L	1558	0	1609	86	0
2	M	1558	0	1609	81	0
2	N	1558	0	1609	75	0
2	V	1558	0	1609	83	0
2	W	1558	0	1609	66	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	X	1558	0	1609	57	0
2	Y	1558	0	1609	74	0
2	Z	1558	0	1609	82	0
All	All	46228	0	47460	2137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 2137 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:12:ILE:HG21	1:D:14:VAL:CG2	1.50	1.41
1:E:12:ILE:HG21	1:E:14:VAL:CG2	1.50	1.40
1:C:12:ILE:HG21	1:C:14:VAL:CG2	1.50	1.39
1:O:12:ILE:HG21	1:O:14:VAL:CG2	1.50	1.39
1:P:12:ILE:HG21	1:P:14:VAL:CG2	1.50	1.39

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2 20
1	B	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2 20
1	C	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2 20
1	D	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2 20
1	E	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2 20
1	F	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2 20
1	G	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2 20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2	20
1	P	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2	20
1	Q	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2	20
1	R	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2	20
1	S	222/233 (95%)	183 (82%)	28 (13%)	11 (5%)	2	20
1	T	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2	20
1	U	222/233 (95%)	184 (83%)	27 (12%)	11 (5%)	2	20
2	1	201/203 (99%)	176 (88%)	23 (11%)	2 (1%)	15	54
2	2	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	H	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	I	201/203 (99%)	183 (91%)	16 (8%)	2 (1%)	15	54
2	J	201/203 (99%)	180 (90%)	20 (10%)	1 (0%)	29	69
2	K	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	L	201/203 (99%)	181 (90%)	16 (8%)	4 (2%)	7	38
2	M	201/203 (99%)	176 (88%)	23 (11%)	2 (1%)	15	54
2	N	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	V	201/203 (99%)	185 (92%)	14 (7%)	2 (1%)	15	54
2	W	201/203 (99%)	183 (91%)	16 (8%)	2 (1%)	15	54
2	X	201/203 (99%)	180 (90%)	20 (10%)	1 (0%)	29	69
2	Y	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	Z	201/203 (99%)	181 (90%)	16 (8%)	4 (2%)	7	38
All	All	5922/6104 (97%)	5120 (86%)	618 (10%)	184 (3%)	7	27

5 of 184 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	12	ILE
1	A	128	GLY
1	A	182	GLU
1	A	200	SER
1	B	12	ILE

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	B	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	C	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	D	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	E	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	F	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	G	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	O	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	P	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	Q	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	R	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	S	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	T	186/193 (96%)	163 (88%)	23 (12%)	4 19
1	U	186/193 (96%)	163 (88%)	23 (12%)	4 19
2	1	170/170 (100%)	152 (89%)	18 (11%)	6 23
2	2	170/170 (100%)	146 (86%)	24 (14%)	3 16
2	H	170/170 (100%)	145 (85%)	25 (15%)	3 15
2	I	170/170 (100%)	148 (87%)	22 (13%)	4 18
2	J	170/170 (100%)	143 (84%)	27 (16%)	2 13
2	K	170/170 (100%)	143 (84%)	27 (16%)	2 13
2	L	170/170 (100%)	146 (86%)	24 (14%)	3 16
2	M	170/170 (100%)	152 (89%)	18 (11%)	6 23
2	N	170/170 (100%)	146 (86%)	24 (14%)	3 16
2	V	170/170 (100%)	145 (85%)	25 (15%)	3 15
2	W	170/170 (100%)	149 (88%)	21 (12%)	4 19
2	X	170/170 (100%)	143 (84%)	27 (16%)	2 13

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	Y	170/170 (100%)	143 (84%)	27 (16%)	2   13
2	Z	170/170 (100%)	146 (86%)	24 (14%)	3   16
All	All	4984/5082 (98%)	4329 (87%)	655 (13%)	7   18

5 of 655 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	T	202	GLU
2	Y	83	LEU
1	U	134	VAL
1	T	201	LEU
2	W	45	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 125 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	164	GLN
2	X	30	ASN
1	P	98	GLN
2	W	164	GLN
2	Z	30	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-1740. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections (i)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices (i)


#### 6.2.1 Primary map



X Index: 72



Y Index: 72




Z Index: 72


The images above show central slices of the map in three orthogonal directions.

### 6.3 Largest variance slices [\(i\)](#)


#### 6.3.1 Primary map



X Index: 90



Y Index: 53




Z Index: 59


The images above show the largest variance slices of the map in three orthogonal directions.

### 6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


#### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [\(i\)](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 2.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.6 Mask visualisation [\(i\)](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis (i)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 295 nm<sup>3</sup>; this corresponds to an approximate mass of 266 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [\(i\)](#)



\*Reported resolution corresponds to spatial frequency of  $0.147 \text{ \AA}^{-1}$


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit i


This section contains information regarding the fit between EMDB map EMD-1740 and PDB model 3C92. Per-residue inclusion information can be found in section 3 on page 6.

### 9.1 Map-model overlay i




The images above show the 3D surface view of the map at the recommended contour level 2.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2.5).

## 9.4 Atom inclusion [\(i\)](#)



At the recommended contour level, 80% of all backbone atoms, 50% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (2.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5030	0.1670
1	0.5230	0.1730
2	0.5270	0.1750
A	0.4800	0.1580
B	0.4860	0.1590
C	0.4900	0.1650
D	0.4810	0.1560
E	0.4920	0.1680
F	0.4980	0.1680
G	0.4860	0.1620
H	0.5220	0.1740
I	0.5280	0.1750
J	0.5300	0.1790
K	0.5220	0.1760
L	0.5250	0.1750
M	0.5280	0.1780
N	0.5280	0.1780
O	0.4590	0.1460
P	0.4730	0.1530
Q	0.4890	0.1660
R	0.4840	0.1620
S	0.4920	0.1660
T	0.4920	0.1660
U	0.4790	0.1570
V	0.5200	0.1710
W	0.5210	0.1670
X	0.5220	0.1700
Y	0.5240	0.1720
Z	0.5220	0.1720

