



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2023 – 06:21 PM EDT

PDB ID : 6D0K
Title : Crystal structure of a CLC-type fluoride/proton antiporter, E118Q mutant
Authors : Last, N.B.; Stockbridge, R.B.; Wilson, A.E.; Shane, T.; Kolmakova-Partensky, L.; Koide, A.; Koide, S.; Miller, C.
Deposited on : 2018-04-10
Resolution : 3.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1558 (3.42-3.30)
Clashscore	141614	1627 (3.42-3.30)
Ramachandran outliers	138981	1599 (3.42-3.30)
Sidechain outliers	138945	1598 (3.42-3.30)
RSRZ outliers	127900	1507 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	
1	B	421	
2	C	93	
2	D	93	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	F	B	503	-	-	X	-
5	MHA	C	101	-	X	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7604 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CLC-type fluoride/proton antiporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	395	3022	2031	468	512	11	0	0	0
1	B	396	3033	2037	472	513	11	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP C9CPP6
A	-1	ALA	-	expression tag	UNP C9CPP6
A	0	ALA	-	expression tag	UNP C9CPP6
A	1	ALA	-	expression tag	UNP C9CPP6
A	4	ILE	MET	engineered mutation	UNP C9CPP6
A	118	GLN	GLU	engineered mutation	UNP C9CPP6
A	407	THR	-	expression tag	UNP C9CPP6
A	408	ARG	-	expression tag	UNP C9CPP6
A	409	GLY	-	expression tag	UNP C9CPP6
A	410	SER	-	expression tag	UNP C9CPP6
A	411	GLY	-	expression tag	UNP C9CPP6
A	412	GLY	-	expression tag	UNP C9CPP6
A	413	HIS	-	expression tag	UNP C9CPP6
A	414	HIS	-	expression tag	UNP C9CPP6
A	415	HIS	-	expression tag	UNP C9CPP6
A	416	HIS	-	expression tag	UNP C9CPP6
A	417	HIS	-	expression tag	UNP C9CPP6
A	418	HIS	-	expression tag	UNP C9CPP6
B	-2	MET	-	expression tag	UNP C9CPP6
B	-1	ALA	-	expression tag	UNP C9CPP6
B	0	ALA	-	expression tag	UNP C9CPP6
B	1	ALA	-	expression tag	UNP C9CPP6
B	4	ILE	MET	engineered mutation	UNP C9CPP6
B	118	GLN	GLU	engineered mutation	UNP C9CPP6
B	407	THR	-	expression tag	UNP C9CPP6

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	408	ARG	-	expression tag	UNP C9CPP6
B	409	GLY	-	expression tag	UNP C9CPP6
B	410	SER	-	expression tag	UNP C9CPP6
B	411	GLY	-	expression tag	UNP C9CPP6
B	412	GLY	-	expression tag	UNP C9CPP6
B	413	HIS	-	expression tag	UNP C9CPP6
B	414	HIS	-	expression tag	UNP C9CPP6
B	415	HIS	-	expression tag	UNP C9CPP6
B	416	HIS	-	expression tag	UNP C9CPP6
B	417	HIS	-	expression tag	UNP C9CPP6
B	418	HIS	-	expression tag	UNP C9CPP6

- Molecule 2 is a protein called Monobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	86	Total 653	C 418	N 100	O 134	S 1	0	0	0
2	C	90	Total 679	C 434	N 104	O 140	S 1	0	0	0

- Molecule 3 is DECYL-BETA-D-MALTOPYRANOSIDE (three-letter code: DMU) (formula: C₂₂H₄₂O₁₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	Total 33	C 22	O 11	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			33	22	11		
3	A	1	Total	C	O	0	0
			33	22	11		
3	A	1	Total	C	O	0	0
			33	22	11		
3	B	1	Total	C	O	0	0
			33	22	11		
3	B	1	Total	C	O	0	0
			26	15	11		

- Molecule 4 is FLUORIDE ION (three-letter code: F) (formula: F).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	F	0	0
			2	2		
4	B	2	Total	F	0	0
			2	2		


- Molecule 5 is (CARBAMOYLMETHYL-CARBOXYMETHYL-AMINO)-ACETIC ACID (three-letter code: MHA) (formula: C₆H₁₀N₂O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			13	6	2	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	3	Total O 3 3	0	0
6	B	6	Total O 6 6	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	118.32Å 125.99Å 134.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.73 – 3.35 49.73 – 3.35	Depositor EDS
% Data completeness (in resolution range)	99.3 (49.73-3.35) 99.7 (49.73-3.35)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 3.33Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.244 , 0.266 0.244 , 0.266	Depositor DCC
R_{free} test set	1442 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	161.2	Xtrriage
Anisotropy	0.496	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 114.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	7604	wwPDB-VP
Average B, all atoms (Å ²)	192.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMU, F, MHA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/3102	0.39	0/4222
1	B	0.25	0/3113	0.39	0/4236
2	C	0.23	0/699	0.44	0/963
2	D	0.23	0/671	0.43	0/921
All	All	0.24	0/7585	0.40	0/10342

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3022	0	3139	19	0
1	B	3033	0	3155	14	0
2	C	679	0	648	4	0
2	D	653	0	618	2	0
3	A	132	0	168	1	0
3	B	59	0	67	0	0
4	A	2	0	0	1	0
4	B	2	0	0	2	0
5	C	13	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	3	0	0	0	0
6	B	6	0	0	0	0
All	All	7604	0	7803	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:118:GLN:N	4:B:503:F:F	2.23	0.61
1:A:101:VAL:HA	1:A:315:LEU:HD13	1.84	0.60
1:A:272:LEU:HD23	1:A:301:LEU:HD13	1.87	0.57
1:B:272:LEU:HD23	1:B:301:LEU:HD13	1.90	0.54
1:A:347:LEU:HD13	1:A:382:VAL:HG21	1.91	0.52
1:B:117:ARG:N	4:B:503:F:F	2.33	0.51
1:B:95:LEU:HD13	1:B:131:ARG:HG3	1.93	0.51
1:A:117:ARG:N	4:A:505:F:F	2.35	0.50
1:B:101:VAL:HA	1:B:315:LEU:HD13	1.94	0.49
1:B:290:THR:HG23	1:B:292:GLN:HG3	1.94	0.48
1:A:104:GLY:HA3	1:A:315:LEU:HD12	1.96	0.47
1:A:58:PRO:HD3	1:A:266:VAL:HG21	1.97	0.47
1:A:95:LEU:HD13	1:A:131:ARG:HG3	1.97	0.47
1:A:276:ARG:HH12	1:A:294:ILE:HG22	1.80	0.46
1:B:347:LEU:HD13	1:B:382:VAL:HG21	1.98	0.46
1:A:290:THR:HG23	1:A:292:GLN:HG3	1.97	0.46
2:C:73:THR:HG23	2:C:88:SER:HB2	1.98	0.45
1:B:286:SER:O	1:B:290:THR:HG22	2.17	0.45
3:A:502:DMU:H8	3:A:502:DMU:H5	1.53	0.45
1:A:144:THR:O	1:A:148:ILE:HG12	2.17	0.45
1:A:276:ARG:NH2	1:A:298:ASP:OD2	2.50	0.44
1:B:97:LEU:HD23	1:B:127:THR:OG1	2.18	0.44
1:B:236:PHE:HD2	1:B:303:LEU:HG	1.82	0.44
2:C:44:ASN:OD1	2:C:44:ASN:N	2.51	0.44
1:B:388:PHE:HA	1:B:391:LYS:HE3	2.00	0.44
1:A:240:LEU:HA	1:A:307:VAL:HG13	2.00	0.43
2:D:44:ASN:OD1	2:D:44:ASN:N	2.51	0.43
1:A:152:ALA:HB2	1:A:188:ALA:HB1	2.01	0.42
1:A:162:ILE:HG23	1:B:380:VAL:HG11	2.00	0.42
1:A:105:THR:OG1	1:A:120:VAL:HG21	2.20	0.42
1:A:388:PHE:HA	1:A:391:LYS:HE3	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:12:VAL:HG22	2:C:22:ILE:HG22	2.03	0.41
2:C:39:GLY:HA3	2:C:47:VAL:HG12	2.02	0.41
1:B:144:THR:O	1:B:148:ILE:HG12	2.20	0.41
1:A:236:PHE:HD2	1:A:303:LEU:HG	1.86	0.41
1:A:271:LEU:HG	2:D:80:LEU:HA	2.03	0.41
1:A:354:GLY:HA3	1:A:363:PRO:HG3	2.02	0.41
1:B:108:THR:O	1:B:113:GLY:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/421 (93%)	379 (96%)	13 (3%)	1 (0%)	41	73
1	B	394/421 (94%)	380 (96%)	13 (3%)	1 (0%)	41	73
2	C	88/93 (95%)	85 (97%)	3 (3%)	0	100	100
2	D	80/93 (86%)	77 (96%)	3 (4%)	0	100	100
All	All	955/1028 (93%)	921 (96%)	32 (3%)	2 (0%)	47	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	THR
1	B	9	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/340 (94%)	320 (100%)	0	100	100
1	B	321/340 (94%)	321 (100%)	0	100	100
2	C	78/80 (98%)	78 (100%)	0	100	100
2	D	74/80 (92%)	74 (100%)	0	100	100
All	All	793/840 (94%)	793 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DMU	A	504	-	34,34,34	0.14	0	45,45,45	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	DMU	B	502	-	27,27,34	0.13	0	37,38,45	0.15	0
3	DMU	B	501	-	34,34,34	0.15	0	45,45,45	0.46	1 (2%)
3	DMU	A	502	-	34,34,34	0.13	0	45,45,45	0.39	0
3	DMU	A	503	-	34,34,34	0.15	0	45,45,45	0.46	0
5	MHA	C	101	-	12,12,12	1.55	4 (33%)	15,15,15	1.44	5 (33%)
3	DMU	A	501	-	34,34,34	0.13	0	45,45,45	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DMU	A	504	-	-	5/19/59/59	0/2/2/2
3	DMU	B	502	-	-	2/12/52/59	0/2/2/2
3	DMU	B	501	-	-	1/19/59/59	0/2/2/2
3	DMU	A	502	-	-	2/19/59/59	0/2/2/2
3	DMU	A	503	-	-	5/19/59/59	0/2/2/2
5	MHA	C	101	-	-	7/12/12/12	-
3	DMU	A	501	-	-	7/19/59/59	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	101	MHA	O2-C2	2.73	1.31	1.22
5	C	101	MHA	O4-C4	2.71	1.31	1.22
5	C	101	MHA	O3-C4	-2.65	1.21	1.30
5	C	101	MHA	O1-C2	-2.62	1.21	1.30

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	101	MHA	C6-C5-N1	-2.92	109.31	114.38
5	C	101	MHA	O3-C4-C3	2.42	123.00	113.45
5	C	101	MHA	O1-C2-C1	2.41	122.95	113.45
5	C	101	MHA	O4-C4-C3	-2.26	113.29	122.31
5	C	101	MHA	O2-C2-C1	-2.23	113.42	122.31
3	B	501	DMU	C6-O5-C4	2.07	117.75	113.69

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	DMU	C19-C18-O16-C6
3	A	502	DMU	C19-C18-O16-C6
3	A	504	DMU	O5-C6-O16-C18
3	B	501	DMU	C19-C18-O16-C6
3	B	502	DMU	C19-C18-O16-C6
5	C	101	MHA	C2-C1-N1-C3
5	C	101	MHA	N1-C1-C2-O1
5	C	101	MHA	N1-C1-C2-O2
5	C	101	MHA	N1-C5-C6-O5
5	C	101	MHA	N1-C5-C6-N2
3	A	501	DMU	O1-C10-O7-C3
3	A	504	DMU	C1-C6-O16-C18
3	A	501	DMU	C1-C6-O16-C18
3	B	502	DMU	O6-C11-C9-O1
3	A	502	DMU	O5-C4-C57-O61
3	A	501	DMU	O6-C11-C9-O1
3	A	503	DMU	O6-C11-C9-O1
3	A	504	DMU	O5-C4-C57-O61
3	A	503	DMU	C19-C18-O16-C6
3	A	501	DMU	C4-C3-O7-C10
3	A	501	DMU	C2-C3-O7-C10
5	C	101	MHA	C2-C1-N1-C5
3	A	501	DMU	O16-C18-C19-C22
3	A	503	DMU	C22-C25-C28-C31
3	A	504	DMU	C2-C3-O7-C10
3	A	503	DMU	C18-C19-C22-C25
3	A	503	DMU	C31-C34-C37-C40
3	A	504	DMU	C4-C3-O7-C10
5	C	101	MHA	N1-C3-C4-O4


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	DMU	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/421 (93%)	0.53	46 (11%) 4 5	148, 178, 229, 272	0
1	B	396/421 (94%)	0.32	31 (7%) 13 15	145, 173, 216, 269	0
2	C	90/93 (96%)	1.86	39 (43%) 0 0	178, 205, 284, 338	0
2	D	86/93 (92%)	1.80	31 (36%) 0 0	198, 248, 304, 338	0
All	All	967/1028 (94%)	0.68	147 (15%) 2 2	145, 181, 257, 338	0

All (147) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	14	ALA	13.9
2	D	12	VAL	13.8
2	C	82	TRP	8.2
2	C	83	TYR	8.1
2	C	45	SER	7.9
2	D	21	LEU	7.7
2	C	42	GLY	7.1
2	D	20	LEU	6.7
2	D	93	THR	6.7
2	C	55	SER	6.5
2	C	40	GLU	6.5
2	D	91	TYR	6.4
2	D	90	ASN	6.3
2	C	56	SER	6.1
2	D	60	THR	5.8
2	C	44	ASN	5.6
1	B	269	PHE	5.6
2	C	53	PRO	5.4
2	D	11	GLU	5.4
2	C	43	GLY	5.1
2	C	81	GLN	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	63	CYS	4.9
2	C	93	THR	4.8
1	B	274	GLN	4.8
1	A	208	LEU	4.7
2	D	15	ALA	4.7
2	D	70	TYR	4.7
1	A	289	PHE	4.7
1	A	91	LYS	4.6
2	C	31	VAL	4.6
1	A	342	GLU	4.6
1	B	52	PRO	4.6
1	A	260	ILE	4.5
2	C	30	SER	4.5
1	B	113	GLY	4.4
1	A	211	SER	4.3
2	C	79	TRP	4.3
1	B	50	PHE	4.2
2	D	59	ALA	4.2
2	D	50	PHE	4.1
2	C	68	VAL	4.0
2	C	39	GLY	4.0
1	B	51	LEU	4.0
2	C	54	GLY	4.0
2	D	92	GLN	3.9
1	A	266	VAL	3.9
2	C	77	HIS	3.9
2	C	80	LEU	3.9
1	A	62	ILE	3.9
1	B	278	SER	3.8
2	C	29	SER	3.8
1	A	267	LEU	3.8
1	A	383	CYS	3.8
1	B	254	PRO	3.8
2	D	48	GLN	3.8
2	C	78	GLY	3.8
2	C	33	TYR	3.7
1	B	275	GLY	3.7
1	A	209	PRO	3.7
1	B	53	LEU	3.7
2	D	68	VAL	3.6
1	A	12	PHE	3.6
1	A	265	LEU	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	67	GLY	3.6
1	A	90	GLU	3.6
2	C	52	VAL	3.5
2	D	22	ILE	3.5
2	C	65	SER	3.5
1	A	247	MET	3.5
1	B	112	GLY	3.5
1	B	265	LEU	3.4
1	B	270	VAL	3.4
1	B	55	LEU	3.4
2	D	66	PRO	3.3
2	C	70	TYR	3.3
2	C	46	PRO	3.3
1	B	253	ASN	3.3
1	A	270	VAL	3.3
1	A	172	LEU	3.1
2	C	32	SER	3.1
1	A	243	MET	3.1
1	A	384	SER	3.0
2	C	34	TYR	3.0
1	A	206	LEU	3.0
2	D	61	ILE	3.0
2	C	84	MET	3.0
2	D	9	LYS	3.0
2	C	66	PRO	3.0
2	D	65	SER	2.9
2	D	49	GLU	2.9
1	B	54	ILE	2.9
2	C	28	SER	2.7
1	B	66	TYR	2.7
1	A	277	TYR	2.7
2	D	83	TYR	2.7
2	D	10	LEU	2.7
1	A	119	GLY	2.7
1	A	51	LEU	2.6
1	B	293	PRO	2.6
1	B	115	ALA	2.6
1	B	316	GLY	2.6
1	A	261	GLY	2.6
2	C	58	THR	2.6
1	A	242	THR	2.6
1	A	387	TYR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	19	LEU	2.5
1	A	386	ALA	2.5
1	A	392	PRO	2.5
1	A	15	THR	2.5
1	A	255	LEU	2.5
1	B	279	GLY	2.5
1	A	50	PHE	2.5
1	B	159	GLU	2.5
1	A	389	ILE	2.5
1	B	114	SER	2.4
2	C	14	ALA	2.4
1	A	390	SER	2.4
2	D	75	TYR	2.4
1	B	276	ARG	2.4
2	D	58	THR	2.4
1	B	315	LEU	2.4
1	B	273	TYR	2.3
2	C	57	SER	2.3
2	C	76	ALA	2.3
1	A	120	VAL	2.3
2	D	38	TYR	2.3
1	A	210	GLN	2.3
2	C	26	ALA	2.3
2	D	69	ASP	2.2
1	A	374	GLN	2.2
1	A	246	ILE	2.2
1	B	264	ALA	2.2
1	A	341	ILE	2.2
2	C	71	THR	2.2
1	A	316	GLY	2.2
2	D	88	SER	2.1
1	B	255	LEU	2.1
1	B	268	LEU	2.1
1	A	59	ILE	2.1
1	B	56	PHE	2.1
2	D	72	ILE	2.1
1	B	260	ILE	2.0
1	A	89	GLU	2.0
1	A	268	LEU	2.0
2	C	41	THR	2.0
1	A	269	PHE	2.0
1	A	375	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	F	A	506	1/1	0.45	0.16	164,164,164,164	0
3	DMU	B	502	26/33	0.79	0.15	212,265,285,289	0
5	MHA	C	101	13/13	0.80	0.11	194,229,240,244	0
3	DMU	A	503	33/33	0.84	0.34	169,227,242,251	0
3	DMU	A	504	33/33	0.85	0.25	156,260,288,292	0
3	DMU	A	501	33/33	0.85	0.18	156,229,258,261	0
3	DMU	A	502	33/33	0.86	0.27	167,214,247,250	0
4	F	B	504	1/1	0.87	0.12	195,195,195,195	0
3	DMU	B	501	33/33	0.88	0.20	156,205,239,245	0
4	F	A	505	1/1	0.96	0.50	166,166,166,166	0
4	F	B	503	1/1	0.97	0.23	164,164,164,164	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DMU B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DMU A 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DMU A 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DMU A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DMU A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DMU B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.