



Full wwPDB NMR Structure Validation Report ⓘ

Mar 1, 2022 – 11:49 AM EST

PDB ID : 2H7T
Title : Solution Structure of the C-terminal Domain of Insulin-like Growth Factor Binding Protein 2 (IGFBP-2)
Authors : Kuang, Z.
Deposited on : 2006-06-03

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.27
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	107	

2 Ensemble composition and analysis

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:187-A:207, A:218-A:271 (75)	1.31	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 9, 10, 11, 12, 13
2	6, 8
3	5, 14
4	18, 19
Single-model clusters	7; 15; 16; 17; 20

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1674 atoms, of which 822 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Insulin-like growth factor-binding protein 2.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	107	1674	523	822	162	158	9	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	183	GLY	-	cloning artifact	UNP P18065

4 Residue-property plots i

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.2 Score per residue for model 2

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.3 Score per residue for model 3

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.4 Score per residue for model 4

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.5 Score per residue for model 5

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.6 Score per residue for model 6

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.7 Score per residue for model 7

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.8 Score per residue for model 8

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.9 Score per residue for model 9

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.10 Score per residue for model 10 (medoid)

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.11 Score per residue for model 11

- Molecule 1: Insulin-like growth factor-binding protein 2



4.2.12 Score per residue for model 12

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.13 Score per residue for model 13

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.14 Score per residue for model 14

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.15 Score per residue for model 15

- Molecule 1: Insulin-like growth factor-binding protein 2



4.2.16 Score per residue for model 16

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.17 Score per residue for model 17

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.18 Score per residue for model 18

- Molecule 1: Insulin-like growth factor-binding protein 2




4.2.19 Score per residue for model 19

- Molecule 1: Insulin-like growth factor-binding protein 2



4.2.20 Score per residue for model 20

- Molecule 1: Insulin-like growth factor-binding protein 2



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.0.3
OPALp	refinement	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.53±0.01	0±0/602 (0.0± 0.0%)	1.00±0.03	0±1/815 (0.0± 0.1%)
All	All	0.53	0/12040 (0.0%)	1.00	4/16300 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.2±1.0
All	All	0	25

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	201	ARG	NE-CZ-NH2	-7.86	116.37	120.30	2	1
1	A	231	TYR	CB-CG-CD2	-6.19	117.28	121.00	19	1
1	A	244	ARG	NE-CZ-NH2	-5.46	117.57	120.30	18	1
1	A	206	ARG	NE-CZ-NH2	-5.28	117.66	120.30	2	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	218	TYR	Sidechain	7
1	A	244	ARG	Sidechain	4
1	A	265	ARG	Sidechain	3
1	A	231	TYR	Sidechain	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	188	ARG	Sidechain	3
1	A	206	ARG	Sidechain	2
1	A	201	ARG	Sidechain	2
1	A	251	ASN	Peptide	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	590	578	578	0±1
All	All	11800	11560	11560	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:233:LEU:HA	1:A:250:VAL:HG22	0.49	1.85	12	1
1:A:240:LEU:HD12	1:A:241:ASN:HD22	0.46	1.69	3	1
1:A:194:GLU:O	1:A:198:VAL:HG23	0.45	2.11	5	2
1:A:228:HIS:H	1:A:228:HIS:CD2	0.43	2.31	7	1
1:A:248:TRP:CD2	1:A:258:ILE:HD12	0.43	2.48	12	1
1:A:261:ALA:HB1	1:A:262:PRO:HD2	0.42	1.91	14	1
1:A:231:TYR:CE2	1:A:255:GLY:HA3	0.41	2.51	19	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	75/107 (70%)	61±3 (82±4%)	11±3 (15±4%)	2±2 (3±2%)	7	39
All	All	1500/2140 (70%)	1228 (82%)	225 (15%)	47 (3%)	7	39

All 22 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	266	GLY	4
1	A	259	GLN	4
1	A	258	ILE	4
1	A	271	HIS	4
1	A	242	GLY	4
1	A	240	LEU	3
1	A	245	GLY	2
1	A	205	MET	2
1	A	251	ASN	2
1	A	262	PRO	2
1	A	268	PRO	2
1	A	265	ARG	2
1	A	239	SER	2
1	A	241	ASN	2
1	A	207	LEU	1
1	A	256	LYS	1
1	A	218	TYR	1
1	A	260	GLY	1
1	A	188	ARG	1
1	A	255	GLY	1
1	A	269	GLU	1
1	A	206	ARG	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	67/95 (71%)	56±4 (84±5%)	11±4 (16±5%)	5	42
All	All	1340/1900 (71%)	1127 (84%)	213 (16%)	5	42

All 52 unique residues with a non-rotameric sidechain are listed below. They are sorted by the

frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	227	LYS	12
1	A	188	ARG	9
1	A	234	LYS	9
1	A	231	TYR	8
1	A	238	MET	8
1	A	254	THR	8
1	A	226	ASP	8
1	A	206	ARG	8
1	A	201	ARG	7
1	A	257	LEU	7
1	A	205	MET	7
1	A	271	HIS	7
1	A	265	ARG	6
1	A	240	LEU	6
1	A	192	GLN	6
1	A	196	ASP	5
1	A	244	ARG	5
1	A	267	ASP	5
1	A	251	ASN	4
1	A	256	LYS	4
1	A	200	GLU	4
1	A	219	SER	4
1	A	236	CYS	4
1	A	233	LEU	4
1	A	232	ASN	4
1	A	197	GLN	3
1	A	203	SER	3
1	A	243	GLN	3
1	A	220	LEU	3
1	A	225	CYS	3
1	A	263	THR	3
1	A	207	LEU	3
1	A	253	ASN	3
1	A	249	CYS	3
1	A	199	LEU	3
1	A	230	LEU	2
1	A	194	GLU	2
1	A	258	ILE	2
1	A	195	LEU	2
1	A	218	TYR	2
1	A	224	ASN	2
1	A	270	CYS	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	237	LYS	1
1	A	241	ASN	1
1	A	189	THR	1
1	A	246	GLU	1
1	A	221	HIS	1
1	A	250	VAL	1
1	A	239	SER	1
1	A	259	GLN	1
1	A	193	GLN	1
1	A	269	GLU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided