



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 05:10 am BST

PDB ID : 4JCJ  
Title : Crystal structure of Isl1 LIM domains with Ldb1 LIM-interaction domain  
Authors : Gadd, M.S.; Jacques, D.A.; Guss, J.M.; Matthews, J.M.  
Deposited on : 2013-02-21  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	173	
1	B	173	
1	C	173	

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 3095 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Insulin gene enhancer protein ISL-1,LIM domain-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	144	991	613	173	189	16	0	0	0
1	B	148	1052	655	185	196	16	0	0	0
1	C	150	1040	644	179	200	17	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	GLY	-	expression tag	UNP P61372
A	10	SER	-	expression tag	UNP P61372
A	288	GLY	-	linker	UNP P61372
A	289	GLY	-	linker	UNP P61372
A	290	SER	-	linker	UNP P61372
A	291	GLY	-	linker	UNP P61372
A	292	GLY	-	linker	UNP P61372
A	293	HIS	-	linker	UNP P61372
A	294	MET	-	linker	UNP P61372
A	295	GLY	-	linker	UNP P61372
A	296	SER	-	linker	UNP P61372
A	297	GLY	-	linker	UNP P61372
A	298	GLY	-	linker	UNP P61372
A	299	GLY	-	linker	UNP P61372
B	9	GLY	-	expression tag	UNP P61372
B	10	SER	-	expression tag	UNP P61372
B	288	GLY	-	linker	UNP P61372
B	289	GLY	-	linker	UNP P61372
B	290	SER	-	linker	UNP P61372
B	291	GLY	-	linker	UNP P61372
B	292	GLY	-	linker	UNP P61372
B	293	HIS	-	linker	UNP P61372

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	294	MET	-	linker	UNP P61372
B	295	GLY	-	linker	UNP P61372
B	296	SER	-	linker	UNP P61372
B	297	GLY	-	linker	UNP P61372
B	298	GLY	-	linker	UNP P61372
B	299	GLY	-	linker	UNP P61372
C	9	GLY	-	expression tag	UNP P61372
C	10	SER	-	expression tag	UNP P61372
C	288	GLY	-	linker	UNP P61372
C	289	GLY	-	linker	UNP P61372
C	290	SER	-	linker	UNP P61372
C	291	GLY	-	linker	UNP P61372
C	292	GLY	-	linker	UNP P61372
C	293	HIS	-	linker	UNP P61372
C	294	MET	-	linker	UNP P61372
C	295	GLY	-	linker	UNP P61372
C	296	SER	-	linker	UNP P61372
C	297	GLY	-	linker	UNP P61372
C	298	GLY	-	linker	UNP P61372
C	299	GLY	-	linker	UNP P61372


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	4	Total Zn 4 4	0	0
2	A	4	Total Zn 4 4	0	0
2	C	4	Total Zn 4 4	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Insulin gene enhancer protein ISL-1,LIM domain-binding protein 1



- Molecule 1: Insulin gene enhancer protein ISL-1,LIM domain-binding protein 1



- Molecule 1: Insulin gene enhancer protein ISL-1,LIM domain-binding protein 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.31Å 56.64Å 179.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.29 – 3.00 48.29 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.8 (48.29-3.00) 98.9 (48.29-3.00)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 3.01Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.254 , 0.299 0.260 , 0.296	Depositor DCC
$R_{free}$ test set	600 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.8	Xtrriage
Anisotropy	0.911	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 81.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.018 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	3095	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/1007	0.62	0/1371
1	B	0.52	0/1070	0.63	0/1454
1	C	0.53	0/1055	0.62	0/1438
All	All	0.54	0/3132	0.62	0/4263

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	991	0	790	16	0
1	B	1052	0	878	30	0
1	C	1040	0	860	28	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
All	All	3095	0	2528	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 72 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:109:ALA:HB3	1:B:132:ASP:OD2	1.74	0.86
1:C:24:ILE:HG21	1:C:29:ILE:HG22	1.68	0.75
1:B:35:ASP:O	1:B:36:LEU:HD12	1.87	0.75
1:B:60:VAL:HG12	1:B:65:THR:HG22	1.69	0.74
1:B:14:ILE:HD11	1:B:16:LEU:HD11	1.68	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/173 (80%)	129 (94%)	9 (6%)	0	100	100
1	B	142/173 (82%)	131 (92%)	9 (6%)	2 (1%)	11	43
1	C	144/173 (83%)	127 (88%)	17 (12%)	0	100	100
All	All	424/519 (82%)	387 (91%)	35 (8%)	2 (0%)	29	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	96	ARG
1	B	68	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/145 (59%)	84 (99%)	1 (1%)	71	90
1	B	95/145 (66%)	90 (95%)	5 (5%)	22	58
1	C	94/145 (65%)	91 (97%)	3 (3%)	39	74
All	All	274/435 (63%)	265 (97%)	9 (3%)	38	73

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	84	ILE
1	C	86	PHE
1	C	55	SER
1	B	57	THR
1	B	86	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	144/173 (83%)	-0.41	0 100 100	31, 54, 80, 91	0
1	B	148/173 (85%)	-0.50	0 100 100	32, 55, 91, 119	0
1	C	150/173 (86%)	-0.49	0 100 100	31, 59, 93, 139	0
All	All	442/519 (85%)	-0.47	0 100 100	31, 56, 89, 139	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	C	403	1/1	0.91	0.04	65,65,65,65	0
2	ZN	C	402	1/1	0.93	0.05	33,33,33,33	0
2	ZN	B	403	1/1	0.93	0.06	41,41,41,41	0
2	ZN	C	404	1/1	0.94	0.03	51,51,51,51	0
2	ZN	A	404	1/1	0.94	0.04	45,45,45,45	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	B	401	1/1	0.96	0.06	43,43,43,43	0
2	ZN	A	402	1/1	0.96	0.07	35,35,35,35	0
2	ZN	B	402	1/1	0.97	0.06	41,41,41,41	0
2	ZN	A	403	1/1	0.98	0.03	46,46,46,46	0
2	ZN	B	404	1/1	0.99	0.03	49,49,49,49	0
2	ZN	A	401	1/1	0.99	0.05	43,43,43,43	0
2	ZN	C	401	1/1	0.99	0.05	31,31,31,31	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.