



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2024 – 11:50 AM EST

PDB ID : 4K4U  
Title : Poliovirus polymerase elongation complex (r5\_form)  
Authors : Gong, P.; Peersen, O.B.  
Deposited on : 2013-04-12  
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)
RNA backbone	3102	1088 (3.12-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	 3% 70% 25%
1	E	471	 3% 73% 21%
2	B	26	 42% 15% 15% 27%
2	F	26	 4% 38% 19% 15% 27%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	16	
3	G	16	
4	D	11	
4	H	11	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9027 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase 3D-POL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	462	3703	2374	611	696	22	0	0	0
1	E	462	3703	2374	611	696	22	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	290	MET	CYS	engineered mutation	UNP P03300
A	446	ASP	LEU	engineered mutation	UNP P03300
A	462	GLY	-	expression tag	UNP P03300
A	463	SER	-	expression tag	UNP P03300
A	464	SER	-	expression tag	UNP P03300
A	465	SER	-	expression tag	UNP P03300
A	466	HIS	-	expression tag	UNP P03300
A	467	HIS	-	expression tag	UNP P03300
A	468	HIS	-	expression tag	UNP P03300
A	469	HIS	-	expression tag	UNP P03300
A	470	HIS	-	expression tag	UNP P03300
A	471	HIS	-	expression tag	UNP P03300
E	290	MET	CYS	engineered mutation	UNP P03300
E	446	ASP	LEU	engineered mutation	UNP P03300
E	462	GLY	-	expression tag	UNP P03300
E	463	SER	-	expression tag	UNP P03300
E	464	SER	-	expression tag	UNP P03300
E	465	SER	-	expression tag	UNP P03300
E	466	HIS	-	expression tag	UNP P03300
E	467	HIS	-	expression tag	UNP P03300
E	468	HIS	-	expression tag	UNP P03300
E	469	HIS	-	expression tag	UNP P03300
E	470	HIS	-	expression tag	UNP P03300
E	471	HIS	-	expression tag	UNP P03300

- Molecule 2 is a RNA chain called RNA (5'-R(\*AP\*AP\*GP\*UP\*CP\*UP\*CP\*CP\*AP\*GP\*GP\*UP\*CP\*UP\*CP\*UP\*CP\*UP\*CP\*UP\*CP\*GP\*UP\*CP\*GP\*AP\*AP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	19	Total	C	N	O	P	0	0	0
			398	178	66	135	19			
2	F	19	Total	C	N	O	P	0	0	0
			398	178	66	135	19			


- Molecule 3 is a RNA chain called RNA (5'-R(\*UP\*GP\*UP\*UP\*CP\*GP\*AP\*CP\*GP\*AP\*GP\*AP\*GP\*AP\*GP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	15	Total	C	N	O	P	0	0	0
			328	146	65	102	15			
3	G	15	Total	C	N	O	P	0	0	0
			328	146	65	102	15			

- Molecule 4 is a RNA chain called RNA (5'-R(P\*GP\*GP\*GP\*GP\*GP\*AP\*GP\*AP\*UP\*GP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	2	Total	C	N	O	P	0	0	0
			46	20	10	14	2			
4	D	2	Total	C	N	O	P	0	0	0
			46	20	10	14	2			

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	25	Total	O	0	0
			25	25		
6	B	3	Total	O	0	0
			3	3		
6	C	3	Total	O	0	0
			3	3		
6	E	16	Total	O	0	0
			16	16		
6	F	4	Total	O	0	0
			4	4		
6	G	6	Total	O	0	0
			6	6		




- Molecule 2: RNA (5'-R(\*AP\*AP\*GP\*UP\*CP\*UP\*CP\*CP\*AP\*GP\*GP\*UP\*CP\*UP\*CP\*UP\*CP\*UP\*CP\*GP\*UP\*CP\*GP\*AP\*AP\*A)-3')

Chain B: 



- Molecule 2: RNA (5'-R(\*AP\*AP\*GP\*UP\*CP\*UP\*CP\*CP\*AP\*GP\*GP\*UP\*CP\*UP\*CP\*UP\*CP\*UP\*CP\*GP\*UP\*CP\*GP\*AP\*AP\*A)-3')

Chain F: 




- Molecule 3: RNA (5'-R(\*UP\*GP\*UP\*UP\*CP\*GP\*AP\*CP\*GP\*AP\*GP\*AP\*GP\*AP\*GP\*A)-3')

Chain C: 



- Molecule 3: RNA (5'-R(\*UP\*GP\*UP\*UP\*CP\*GP\*AP\*CP\*GP\*AP\*GP\*AP\*GP\*AP\*GP\*A)-3')

Chain G: 



- Molecule 4: RNA (5'-R(P\*GP\*GP\*GP\*GP\*GP\*AP\*GP\*AP\*UP\*GP\*A)-3')

Chain H: 



- Molecule 4: RNA (5'-R(P\*GP\*GP\*GP\*GP\*GP\*AP\*GP\*AP\*UP\*GP\*A)-3')

Chain D: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.65Å 63.87Å 105.09Å 101.08° 106.22° 103.03°	Depositor
Resolution (Å)	43.26 – 2.85 43.26 – 2.82	Depositor EDS
% Data completeness (in resolution range)	98.0 (43.26-2.85) 97.4 (43.26-2.82)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.72 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, $R_{free}$	0.219 , 0.276 0.219 , 0.276	Depositor DCC
$R_{free}$ test set	1802 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	83.2	Xtrriage
Anisotropy	0.556	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 56.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.54$ , $\langle L^2 \rangle = 0.39$	Xtrriage
Estimated twinning fraction	0.439 for k,h,-h-k-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9027	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	96.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.48	0/3793	0.63	0/5129
1	E	0.48	0/3793	0.63	0/5129
2	B	0.75	0/442	1.33	3/685 (0.4%)
2	F	0.68	0/442	1.21	3/685 (0.4%)
3	C	0.63	0/368	1.18	1/573 (0.2%)
3	G	0.69	0/368	1.18	1/573 (0.2%)
4	D	0.35	0/51	1.01	0/78
4	H	0.33	0/51	0.93	0/78
All	All	0.52	0/9308	0.79	8/12930 (0.1%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	606	C	N3-C4-C5	-7.89	118.74	121.90
3	C	697	A	O5'-P-OP2	-6.72	99.65	105.70
2	F	606	C	N3-C4-C5	-6.48	119.31	121.90
2	F	606	C	N3-C4-N4	6.23	122.36	118.00
2	B	606	C	N3-C4-N4	5.40	121.78	118.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3703	0	3666	73	0
1	E	3703	0	3666	67	0
2	B	398	0	205	6	0
2	F	398	0	205	9	0
3	C	328	0	164	5	0
3	G	328	0	164	4	0
4	D	46	0	23	1	0
4	H	46	0	23	1	0
5	A	10	0	0	2	0
5	E	10	0	0	1	0
6	A	25	0	0	1	0
6	B	3	0	0	0	0
6	C	3	0	0	0	0
6	E	16	0	0	3	0
6	F	4	0	0	0	0
6	G	6	0	0	0	0
All	All	9027	0	8116	154	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 154 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:ASP:H	1:A:217:PHE:HD2	1.31	0.76
1:A:290:MET:HB2	1:A:293:THR:HG23	1.73	0.69
1:E:269:ASN:N	1:E:269:ASN:OD1	2.23	0.67
1:E:29:ALA:HB3	1:E:441:ILE:HD11	1.76	0.67
1:E:433:LEU:HD13	1:E:437:ARG:HH12	1.59	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	460/471 (98%)	411 (89%)	41 (9%)	8 (2%)	9	27
1	E	460/471 (98%)	411 (89%)	43 (9%)	6 (1%)	12	33
All	All	920/942 (98%)	822 (89%)	84 (9%)	14 (2%)	10	30

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	LYS
1	A	227	GLU
1	E	139	LYS
1	E	227	GLU
1	A	33	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	403/412 (98%)	367 (91%)	36 (9%)	9	26
1	E	403/412 (98%)	369 (92%)	34 (8%)	11	28
All	All	806/824 (98%)	736 (91%)	70 (9%)	10	27

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	260	ASP
1	E	269	ASN
1	E	396	GLU
1	A	294	SER
1	A	269	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	18/26 (69%)	4 (22%)	0
2	F	18/26 (69%)	4 (22%)	0
3	C	14/16 (87%)	2 (14%)	0
3	G	14/16 (87%)	2 (14%)	0
4	D	1/11 (9%)	0	0
4	H	1/11 (9%)	0	0
All	All	66/106 (62%)	12 (18%)	0

5 of 12 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	602	C
2	B	611	C
2	B	613	A
2	B	614	A
3	C	688	U

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	SO4	E	501	-	4,4,4	0.24	0	6,6,6	0.09	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	SO4	A	502	-	4,4,4	0.27	0	6,6,6	0.23	0
5	SO4	E	502	-	4,4,4	0.15	0	6,6,6	0.28	0
5	SO4	A	501	-	4,4,4	0.20	0	6,6,6	0.24	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	502	SO4	1	0
5	E	502	SO4	1	0
5	A	501	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	462/471 (98%)	0.07	13 (2%) 53 48	64, 92, 116, 132	0
1	E	462/471 (98%)	0.14	12 (2%) 56 52	66, 91, 116, 132	0
2	B	19/26 (73%)	-0.06	0 100 100	73, 105, 177, 198	0
2	F	19/26 (73%)	0.07	1 (5%) 26 22	75, 105, 177, 198	0
3	C	15/16 (93%)	-0.33	0 100 100	71, 97, 158, 161	0
3	G	15/16 (93%)	-0.30	0 100 100	71, 97, 158, 162	0
4	D	2/11 (18%)	1.43	1 (50%) 0 0	170, 170, 170, 185	0
4	H	2/11 (18%)	0.19	0 100 100	170, 170, 170, 184	0
All	All	996/1048 (95%)	0.09	27 (2%) 54 50	64, 92, 121, 198	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	321	LEU	4.6
1	A	312	THR	4.4
1	A	313	TYR	4.0
1	A	321	LEU	3.9
1	E	312	THR	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	E	502	5/5	0.82	0.12	118,123,128,138	0
5	SO4	A	501	5/5	0.87	0.14	104,114,120,133	0
5	SO4	A	502	5/5	0.90	0.09	121,123,126,140	0
5	SO4	E	501	5/5	0.94	0.07	103,114,123,131	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.