Full wwPDB NMR Structure Validation Report (i)

Apr 21, 2024 – 05:12 AM EDT

PDB ID	:	2LTK
BMRB ID	:	18485
Title	:	Solution structure of a monomeric truncated mutant of Trypanosoma brucei 1-C-Grx1
Authors	:	Pavan, C.; Gesiot, L.; Sturlese, M.; Mammi, S.; Manta, B.; Comini, M.; Bellanda, M.
Deposited on	:	2012-05-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 94%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	110	91% 6% •

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 14 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:78-A:184 (107)	0.48	14

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	3, 4, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 20
2	1, 2, 10, 18
3	5, 17, 19

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1755 atoms, of which 894 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Mono-cysteine glutaredoxin.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	110	1755	554	894	137	163	7	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	75	GLY	-	expression tag	UNP Q2UZM9

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Mono-cysteine glutaredoxin

Chain A:

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: Mono-cysteine glutaredoxin

Chain A:



4.2.2 Score per residue for model 2

• Molecule 1: Mono-cysteine glutaredoxin

Chain A:



4.2.3 Score per residue for model 3

• Molecule 1: Mono-cysteine glutaredoxin

Chain A: 86% 9%

4.2.4 Score per residue for model 4

• Molecule 1: Mono-cysteine glutaredoxin

Chain A: 87% 10%

4.2.5 Score per residue for model 5

• Molecule 1: Mono-cysteine glutaredoxin

Chain A: 87% 10%

4.2.6 Score per residue for model 6

• Molecule 1: Mono-cysteine glutaredoxin

Chain A: 87% 10%

4.2.7 Score per residue for model 7

• Molecule 1: Mono-cysteine glutaredoxin

Chain A: 89% 8%

4.2.8 Score per residue for model 8

- Molecule 1: Mono-cysteine glutaredoxin

Chain A: 85% 13%

4.2.9 Score per residue for model 9

- Molecule 1: Mono-cysteine glutaredoxin

Chain A: 88% 8%

4.2.10 Score per residue for model 10

- Molecule 1: Mono-cysteine glutaredoxin

Chain A: 88% 9%

4.2.11 Score per residue for model 11

- Molecule 1: Mono-cysteine glutaredoxin

Chain A: 90% 6%

4.2.12 Score per residue for model 12

- Molecule 1: Mono-cysteine glutaredoxin

Chain A: 83% 15%



4.2.13 Score per residue for model 13

• Molecule 1: Mono-cysteine glutaredoxin

Chain A:



4.2.14 Score per residue for model 14 (medoid)

• Molecule 1: Mono-cysteine glutaredoxin

Chain A:

4.2.15 Score per residue for model 15

• Molecule 1: Mono-cysteine glutaredoxin

Chain A:



4.2.16 Score per residue for model 16

• Molecule 1: Mono-cysteine glutaredoxin

Chain A:

4.2.17 Score per residue for model 17

• Molecule 1: Mono-cysteine glutaredoxin

Chain A:

4.2.18 Score per residue for model 18

- Molecule 1: Mono-cysteine glutaredoxin

Chain A:



4.2.19 Score per residue for model 19

- Molecule 1: Mono-cysteine glutaredoxin

Chain A:



4.2.20 Score per residue for model 20

- Molecule 1: Mono-cysteine glutaredoxin

Chain A:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
Amber	refinement	
UNIO'10	structure solution	2.0.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1415
Number of shifts mapped to atoms	1415
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	94%

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.67±0.00	0±0/858 (0.0± 0.0%)	1.05±0.02	3±1/1156 (0.3± 0.1%)
All	All	0.67	0/17160 (0.0%)	1.05	63/23120 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.3±0.5
All	All	0	7

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	109	ARG	NE-CZ-NH1	9.29	124.95	120.30	10	15
1	A	133	ARG	NE-CZ-NH1	9.04	124.82	120.30	1	15
1	A	175	ARG	NE-CZ-NH1	9.03	124.81	120.30	4	17
1	A	182	ARG	NE-CZ-NH1	8.60	124.60	120.30	20	14
1	A	121	TYR	CB-CG-CD1	-5.29	117.83	121.00	3	1
1	A	125	ASP	CB-CG-OD1	5.14	122.92	118.30	9	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	182	ARG	Peptide	2
1	A	135	TYR	Sidechain	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	97	GLY	Peptide	1
1	A	121	TYR	Sidechain	1
1	A	109	ARG	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	16880	17500	17500	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	106/110 (96%)	103±1 (97±1%)	3±1 (3±1%)	0±0 (0±0%)	54 85
All	All	2120/2200 (96%)	2057 (97%)	61 (3%)	2 (0%)	54 85

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	141	GLU	2

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/97 (99%)	89±2 (92±2%)	7±2 (8±2%)	16	64
All	All	1920/1940 (99%)	1771 (92%)	149 (8%)	16	64

All 30 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	173	MET	20
1	A	98	LEU	17
1	A	163	LYS	16
1	A	120	GLU	10
1	A	148	LEU	10
1	A	176	ASP	9
1	A	125	ASP	7
1	A	170	LEU	7
1	A	108	LYS	6
1	A	109	ARG	5
1	A	79	LYS	4
1	A	110	MET	4
1	A	129	HIS	4
1	A	87	LYS	4
1	A	137	LYS	3
1	A	177	LYS	3
1	A	80	LYS	3
1	A	131	VAL	2
1	A	172	LYS	2
1	A	183	ASP	2
1	A	134	SER	2
1	A	145	ILE	1
1	A	158	LEU	1
1	A	169	ASP	1
1	A	104	CYS	1
1	A	184	LEU	1
1	A	96	LYS	1
1	A	180	THR	1
1	A	159	ASP	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	165	LEU	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 94% for the well-defined parts and 93% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1415
Number of shifts mapped to atoms	1415
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	109	-0.20 ± 0.11	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	103	0.14 ± 0.16	None needed (< 0.5 ppm)
$^{13}C'$	107	0.01 ± 0.09	None needed (< 0.5 ppm)
^{15}N	101	0.78 ± 0.20	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 94%, i.e. 1393 atoms were assigned a chemical shift out of a possible 1487. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	526/531 (99%)	214/215 (100%)	212/214 (99%)	100/102 (98%)
Sidechain	801/870 (92%)	549/569 (96%)	251/277 (91%)	1/24 (4%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	66/86 (77%)	40/42 (95%)	25/42 (60%)	1/2 (50%)
Overall	1393/1487 (94%)	803/826 (97%)	488/533 (92%)	102/128 (80%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 93%, i.e. 1415 atoms were assigned a chemical shift out of a possible 1517. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	534/547 (98%)	217/222 (98%)	216/220 (98%)	101/105 (96%)
Sidechain	815/884 (92%)	559/579 (97%)	255/281 (91%)	1/24 (4%)
Aromatic	66/86 (77%)	40/42 (95%)	25/42 (60%)	1/2 (50%)
Overall	1415/1517 (93%)	816/843 (97%)	496/543 (91%)	103/131 (79%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	146	PRO	CB	37.66	26.06 – 37.61	5.0

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:


RCI
0.00
0.02
0.04
0.06
0.08
0.10
76:ALA
80:LYS
84:ASP
88:SER
92:VAL
96:LYS
100:GLU
104:CYS
108:LYS
112:ASP
116:ALA
120:GLU
124:PHE
128:ALA
132:VAL
136:VAL
140:SER
144:THR
148:LEU
152:ALA
156:GLY
160:ILE
164:MET
168:GLY
172:LYS
176:ASP
180:THR
184:LEU
Auth_seq_ID