



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 09:30 am BST

PDB ID : 5NVY  
Title : pVHL:EloB:EloC in complex with (2S,4R)-1-((S)-2-acetamidopropanoyl)-4-hydroxy-N-(4-(4-methylthiazol-5-yl)benzyl) pyrrolidine-2-carboxamide (ligand 11)  
Authors : Soares, P.; Gadd, M.S.; Ciulli, A.  
Deposited on : 2017-05-04  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	
1	D	104	
1	G	104	
1	J	104	
2	B	97	
2	E	97	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	97	 <p>2% 91% 8%</p>
2	K	97	 <p>85% 6% 9%</p>
3	C	162	 <p>% 84% 13%</p>
3	F	162	 <p>3% 82% 6% 12%</p>
3	I	162	 <p>2% 83% 6% 12%</p>
3	L	162	 <p>2% 86% 12%</p>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 11274 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Elongin-B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	As	C	N	O	S			
1	A	104	829	2	524	138	160	5	0	0	0
1	D	103	819	2	518	136	158	5	0	0	0
1	G	103	819	2	518	136	158	5	0	0	0
1	J	103	819	2	518	136	158	5	0	0	0

- Molecule 2 is a protein called Elongin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	87	694	448	111	128	7	0	0	0
2	E	91	723	463	116	137	7	0	0	0
2	H	89	706	454	113	132	7	0	0	0
2	K	88	701	452	112	130	7	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	16	MET	-	initiating methionine	UNP Q15369
E	16	MET	-	initiating methionine	UNP Q15369
H	16	MET	-	initiating methionine	UNP Q15369
K	16	MET	-	initiating methionine	UNP Q15369

- Molecule 3 is a protein called Von Hippel-Lindau disease tumor suppressor.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	S	0	0
			30	21	4	4	1		
4	F	1	Total	C	N	O	S	0	0
			30	21	4	4	1		
4	I	1	Total	C	N	O	S	0	0
			30	21	4	4	1		
4	L	1	Total	C	N	O	S	0	0
			30	21	4	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	51	Total	O	0	0
			51	51		
5	B	18	Total	O	0	0
			18	18		
5	C	38	Total	O	0	0
			38	38		
5	D	20	Total	O	0	0
			20	20		
5	E	21	Total	O	0	0
			21	21		
5	F	35	Total	O	0	0
			35	35		
5	G	31	Total	O	0	0
			31	31		
5	H	21	Total	O	0	0
			21	21		
5	I	33	Total	O	0	0
			33	33		
5	J	38	Total	O	0	0
			38	38		
5	K	25	Total	O	0	0
			25	25		
5	L	35	Total	O	0	0
			35	35		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Elongin-B




- Molecule 1: Elongin-B




- Molecule 1: Elongin-B




- Molecule 1: Elongin-B



- Molecule 2: Elongin-C




- Molecule 2: Elongin-C




- Molecule 2: Elongin-C




- Molecule 2: Elongin-C




- Molecule 3: Von Hippel-Lindau disease tumor suppressor




- Molecule 3: Von Hippel-Lindau disease tumor suppressor



- Molecule 3: Von Hippel-Lindau disease tumor suppressor



- Molecule 3: Von Hippel-Lindau disease tumor suppressor



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.08Å 93.08Å 362.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.76 – 2.90 48.76 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.76-2.90) 99.9 (48.76-2.90)	Depositor EDS
$R_{merge}$	0.23	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.48 (at 2.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0123	Depositor
R, $R_{free}$	0.203 , 0.269 0.204 , 0.269	Depositor DCC
$R_{free}$ test set	1893 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.9	Xtrriage
Anisotropy	0.788	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 29.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11274	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.38 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5666e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, 9B5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/825	0.65	0/1110
1	D	0.39	0/815	0.60	0/1099
1	G	0.40	0/815	0.60	0/1099
1	J	0.38	0/815	0.62	0/1099
2	B	0.38	0/708	0.55	0/954
2	E	0.39	0/737	0.53	0/993
2	H	0.37	0/720	0.53	0/970
2	K	0.37	0/715	0.60	0/964
3	C	0.39	0/1176	0.69	0/1604
3	F	0.40	0/1194	0.66	0/1628
3	I	0.45	1/1194 (0.1%)	0.66	0/1628
3	L	0.40	0/1194	0.65	0/1628
All	All	0.40	1/10908 (0.0%)	0.62	0/14776

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	204	GLU	CD-OE2	7.03	1.33	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	829	0	821	4	0
1	D	819	0	808	1	0
1	G	819	0	808	4	0
1	J	819	0	808	1	0
2	B	694	0	694	2	0
2	E	723	0	719	3	0
2	H	706	0	707	1	0
2	K	701	0	701	2	0
3	C	1156	0	1152	4	0
3	F	1174	0	1166	5	0
3	I	1174	0	1166	3	0
3	L	1174	0	1166	0	0
4	C	30	0	0	0	0
4	F	30	0	0	0	0
4	I	30	0	0	1	0
4	L	30	0	0	0	0
5	A	51	0	0	1	0
5	B	18	0	0	0	0
5	C	38	0	0	0	0
5	D	20	0	0	0	0
5	E	21	0	0	0	0
5	F	35	0	0	0	0
5	G	31	0	0	0	0
5	H	21	0	0	1	0
5	I	33	0	0	0	0
5	J	38	0	0	0	0
5	K	25	0	0	0	0
5	L	35	0	0	0	0
All	All	11274	0	10716	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:102:VAL:HG11	3:F:170:VAL:HG13	1.78	0.66
1:G:8:ARG:HG2	1:G:13:THR:HG23	1.86	0.58
2:E:62:PHE:CD1	2:E:65:ILE:HD12	2.43	0.53
1:G:7:ILE:HD11	1:G:27:LEU:HD11	1.93	0.51
2:E:95:ILE:HB	3:F:165:VAL:HG21	1.94	0.50
3:C:180:ILE:HD12	3:C:184:LEU:HB3	1.92	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:107:ARG:NH1	4:I:301:9B5:NAP	2.61	0.48
1:A:8:ARG:HG2	1:A:13:THR:HG23	1.94	0.48
1:A:62:PHE:CZ	1:A:75:VAL:HG22	2.50	0.46
2:B:101:LEU:HD11	3:C:178:LEU:HD22	1.97	0.46
3:F:180:ILE:HD12	3:F:184:LEU:HB3	1.98	0.46
2:E:37:LEU:HD22	2:E:43:LYS:HG3	1.98	0.45
3:C:180:ILE:HD12	3:C:184:LEU:CB	2.47	0.45
2:K:38:THR:HG23	2:K:80:LYS:HD3	1.98	0.45
3:C:120:ARG:NH1	3:C:197:ASP:OD2	2.50	0.44
1:G:102:VAL:HG11	3:I:174:ASN:HB3	2.00	0.44
1:G:37:ARG:NH1	1:G:41:GLU:OE1	2.51	0.44
3:I:84:VAL:HG22	3:I:128:LEU:CD1	2.48	0.44
1:J:62:PHE:CZ	1:J:75:VAL:HG22	2.53	0.43
1:A:88:LEU:HD21	5:A:212:HOH:O	2.19	0.42
2:B:69:VAL:HG21	2:B:102:GLU:HB3	2.01	0.42
3:F:62:VAL:HG11	3:F:202:THR:HG23	2.00	0.42
2:H:82:ARG:NE	5:H:201:HOH:O	2.51	0.42
2:K:39:SER:HB3	2:K:42:ILE:HD12	2.01	0.41
1:A:43:ARG:HG2	1:A:50:LEU:HD13	2.03	0.40
3:F:141:ASN:HD22	3:F:141:ASN:N	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/104 (96%)	96 (96%)	3 (3%)	1 (1%)	15	45
1	D	99/104 (95%)	95 (96%)	4 (4%)	0	100	100
1	G	99/104 (95%)	95 (96%)	4 (4%)	0	100	100
1	J	99/104 (95%)	93 (94%)	4 (4%)	2 (2%)	7	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	83/97 (86%)	82 (99%)	1 (1%)	0	100	100
2	E	87/97 (90%)	87 (100%)	0	0	100	100
2	H	85/97 (88%)	83 (98%)	2 (2%)	0	100	100
2	K	84/97 (87%)	81 (96%)	3 (4%)	0	100	100
3	C	138/162 (85%)	134 (97%)	4 (3%)	0	100	100
3	F	140/162 (86%)	136 (97%)	4 (3%)	0	100	100
3	I	140/162 (86%)	134 (96%)	6 (4%)	0	100	100
3	L	140/162 (86%)	134 (96%)	6 (4%)	0	100	100
All	All	1294/1452 (89%)	1250 (97%)	41 (3%)	3 (0%)	47	78

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	ASP
1	J	82	ASP
1	J	83	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/90 (100%)	88 (98%)	2 (2%)	52	81
1	D	89/90 (99%)	87 (98%)	2 (2%)	52	81
1	G	89/90 (99%)	87 (98%)	2 (2%)	52	81
1	J	89/90 (99%)	88 (99%)	1 (1%)	73	92
2	B	78/86 (91%)	75 (96%)	3 (4%)	33	67
2	E	82/86 (95%)	80 (98%)	2 (2%)	49	79
2	H	80/86 (93%)	80 (100%)	0	100	100
2	K	79/86 (92%)	77 (98%)	2 (2%)	47	78
3	C	131/147 (89%)	131 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	F	133/147 (90%)	130 (98%)	3 (2%)	50	80
3	I	133/147 (90%)	129 (97%)	4 (3%)	41	75
3	L	133/147 (90%)	130 (98%)	3 (2%)	50	80
All	All	1206/1292 (93%)	1182 (98%)	24 (2%)	55	82

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	ASP
1	A	83	ASP
2	B	63	ARG
2	B	72	LYS
2	B	101	LEU
1	D	9	ARG
1	D	86	GLU
2	E	17	MET
2	E	56	GLU
3	F	69	ARG
3	F	73	GLN
3	F	203	GLN
1	G	12	THR
1	G	83	ASP
3	I	148	PHE
3	I	161	ARG
3	I	180	ILE
3	I	182	ARG
1	J	52	ASP
2	K	85	ASN
2	K	101	LEU
3	L	69	ARG
3	L	148	PHE
3	L	161	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	F	141	ASN
3	F	203	GLN
1	J	10	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	CAS	I	77	3	5,8,9	1.00	0	1,9,11	0.59	0
1	CAS	G	89	1	5,8,9	1.12	0	1,9,11	0.54	0
3	CAS	C	77	3	5,8,9	0.76	0	1,9,11	0.40	0
1	CAS	A	60	1	5,8,9	0.87	0	1,9,11	0.72	0
3	CAS	F	77	3	5,8,9	0.68	0	1,9,11	0.01	0
1	CAS	G	60	1	5,8,9	0.90	0	1,9,11	0.47	0
1	CAS	D	60	1	5,8,9	0.90	0	1,9,11	0.43	0
1	CAS	A	89	1	5,8,9	1.08	0	1,9,11	0.58	0
1	CAS	D	89	1	5,8,9	1.07	0	1,9,11	0.32	0
1	CAS	J	60	1	5,8,9	0.97	0	1,9,11	0.38	0
1	CAS	J	89	1	5,8,9	1.08	0	1,9,11	0.15	0
3	CAS	L	77	3	5,8,9	0.95	0	1,9,11	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CAS	I	77	3	-	0/0/7/9	-
1	CAS	G	89	1	-	0/0/7/9	-
3	CAS	C	77	3	-	0/0/7/9	-
1	CAS	A	60	1	-	0/0/7/9	-
3	CAS	F	77	3	-	0/0/7/9	-
1	CAS	G	60	1	-	0/0/7/9	-
1	CAS	D	60	1	-	0/0/7/9	-
1	CAS	A	89	1	-	0/0/7/9	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	D	89	1	-	0/0/7/9	-
1	CAS	J	60	1	-	0/0/7/9	-
1	CAS	J	89	1	-	0/0/7/9	-
3	CAS	L	77	3	-	0/0/7/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	9B5	F	301	-	27,32,32	2.94	6 (22%)	34,45,45	1.33	5 (14%)
4	9B5	C	301	-	27,32,32	3.03	5 (18%)	34,45,45	1.18	3 (8%)
4	9B5	L	301	-	27,32,32	2.99	6 (22%)	34,45,45	1.23	3 (8%)
4	9B5	I	301	-	27,32,32	3.04	6 (22%)	34,45,45	1.25	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	9B5	F	301	-	-	0/25/37/37	0/3/3/3
4	9B5	C	301	-	-	0/25/37/37	0/3/3/3
4	9B5	L	301	-	-	0/25/37/37	0/3/3/3
4	9B5	I	301	-	-	0/25/37/37	0/3/3/3

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	I	301	9B5	CAY-CAZ	-13.50	1.35	1.48
4	C	301	9B5	CAY-CAZ	-13.30	1.35	1.48
4	L	301	9B5	CAY-CAZ	-12.93	1.35	1.48
4	F	301	9B5	CAY-CAZ	-12.91	1.35	1.48
4	L	301	9B5	CBC-CAU	-4.97	1.41	1.52
4	I	301	9B5	CBC-CAU	-4.75	1.41	1.52
4	C	301	9B5	CBC-CAU	-4.75	1.41	1.52
4	F	301	9B5	CBC-CAU	-4.54	1.42	1.52
4	I	301	9B5	CAM-CAW	-4.32	1.42	1.51
4	L	301	9B5	CAM-CAW	-4.31	1.42	1.51
4	C	301	9B5	CAM-CAW	-4.27	1.42	1.51
4	F	301	9B5	CAM-CAW	-4.18	1.42	1.51
4	C	301	9B5	CBC-NBD	3.22	1.53	1.47
4	F	301	9B5	CBC-NBD	3.04	1.53	1.47
4	L	301	9B5	CBC-NBD	2.64	1.52	1.47
4	L	301	9B5	CAO-NBD	2.62	1.51	1.47
4	I	301	9B5	CAO-NBD	2.54	1.51	1.47
4	I	301	9B5	CBC-NBD	2.28	1.51	1.47
4	C	301	9B5	CAB-CAX	2.23	1.54	1.50
4	I	301	9B5	CAB-CAX	2.15	1.54	1.50
4	F	301	9B5	CAO-NBD	2.11	1.50	1.47
4	L	301	9B5	CAB-CAX	2.09	1.53	1.50
4	F	301	9B5	CAB-CAX	2.09	1.53	1.50

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	301	9B5	CBA-CAO-NBD	4.47	107.83	103.08
4	I	301	9B5	CBA-CAO-NBD	3.91	107.24	103.08
4	L	301	9B5	CBA-CAO-NBD	3.33	106.62	103.08
4	C	301	9B5	CBA-CAO-NBD	3.24	106.53	103.08
4	F	301	9B5	CBA-CAN-CBC	2.97	107.77	103.97
4	L	301	9B5	CAO-NBD-CBC	-2.49	108.11	111.70
4	F	301	9B5	CAN-CBA-CAO	2.48	106.35	103.31
4	C	301	9B5	CAO-NBD-CBC	-2.44	108.18	111.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	I	301	9B5	CBA-CAN-CBC	2.43	107.08	103.97
4	I	301	9B5	CAW-CAM-NAQ	-2.42	107.87	113.05
4	I	301	9B5	CAO-NBD-CBC	-2.33	108.33	111.70
4	L	301	9B5	CBA-CAN-CBC	2.27	106.87	103.97
4	C	301	9B5	CAN-CBC-CAU	-2.22	106.89	111.32
4	F	301	9B5	CAO-NBD-CBC	-2.15	108.59	111.70
4	F	301	9B5	CAN-CBC-NBD	2.03	105.78	103.10

There are no chirality outliers.



There are no torsion outliers.



There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	I	301	9B5	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	102/104 (98%)	-0.23	1 (0%) 82 82	34, 50, 92, 117	0
1	D	101/104 (97%)	-0.10	3 (2%) 50 45	44, 69, 110, 150	0
1	G	101/104 (97%)	-0.15	1 (0%) 82 82	48, 67, 95, 109	0
1	J	101/104 (97%)	-0.20	1 (0%) 82 82	36, 50, 95, 132	0
2	B	87/97 (89%)	-0.33	0 100 100	33, 49, 80, 102	0
2	E	91/97 (93%)	-0.18	2 (2%) 62 59	41, 60, 103, 114	0
2	H	89/97 (91%)	-0.01	2 (2%) 62 59	48, 66, 97, 110	0
2	K	88/97 (90%)	-0.25	0 100 100	37, 52, 83, 96	0
3	C	140/162 (86%)	-0.11	2 (1%) 75 75	37, 55, 94, 114	0
3	F	142/162 (87%)	-0.00	5 (3%) 44 38	41, 61, 97, 113	0
3	I	142/162 (87%)	-0.13	3 (2%) 63 61	40, 56, 115, 132	0
3	L	142/162 (87%)	-0.18	3 (2%) 63 61	32, 48, 109, 144	0
All	All	1326/1452 (91%)	-0.15	23 (1%) 70 69	32, 57, 101, 150	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	101	ASP	4.9
3	L	143	ASP	4.0
3	I	186	GLU	3.0
3	F	179	ASP	3.0
3	C	143	ASP	3.0
3	F	143	ASP	2.9
2	H	109	PHE	2.9
1	D	81	ALA	2.8
1	D	83	ASP	2.7
3	F	177	ARG	2.5
3	F	185	TYR	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	E	56	GLU	2.4
3	C	185	TYR	2.4
2	E	55	ASN	2.3
3	I	177	ARG	2.3
1	D	84	THR	2.2
3	L	181	VAL	2.2
2	H	85	ASN	2.1
1	G	1	MET	2.1
3	L	182	ARG	2.1
3	I	184	LEU	2.0
3	F	139	SER	2.0
1	A	101	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CAS	A	60	9/10	0.84	0.23	49,52,84,92	0
1	CAS	J	60	9/10	0.90	0.24	51,56,89,93	0
1	CAS	G	60	9/10	0.92	0.15	61,64,86,86	0
1	CAS	D	60	9/10	0.92	0.14	53,57,81,82	0
1	CAS	D	89	9/10	0.92	0.18	93,99,153,155	0
1	CAS	G	89	9/10	0.92	0.14	66,73,110,114	0
1	CAS	A	89	9/10	0.95	0.15	61,68,116,116	0
1	CAS	J	89	9/10	0.95	0.12	54,58,89,94	0
3	CAS	I	77	9/10	0.96	0.12	45,47,66,67	0
3	CAS	C	77	9/10	0.97	0.13	49,52,71,74	0
3	CAS	L	77	9/10	0.97	0.11	39,43,64,64	0
3	CAS	F	77	9/10	0.98	0.12	56,60,81,85	0

## 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	9B5	F	301	30/30	0.93	0.23	32,48,55,58	0
4	9B5	L	301	30/30	0.95	0.19	25,32,50,67	0
4	9B5	I	301	30/30	0.95	0.19	27,38,50,57	0
4	9B5	C	301	30/30	0.96	0.19	27,39,49,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around 9B5 L 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 9B5 I 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.