



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 3, 2023 – 06:15 am GMT

PDB ID : 1ODK
Title : PURINE NUCLEOSIDE PHOSPHORYLASE FROM THERMUS THERMOPHILUS
Authors : Tahirov, T.H.; Inagaki, E.; Miyano, M.
Deposited on : 2003-02-19
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

PERCENTILES INFOmissingINFO

1 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 11369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PURINE NUCLEOSIDE PHOSPHORYLASE.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 1 | A | 228 | Total 1745 | C 1108 | N 319 | O 314 | S 4 | 0 | 0 | 0 |
| 1 | B | 234 | Total 1782 | C 1131 | N 325 | O 322 | S 4 | 0 | 0 | 0 |
| 1 | C | 228 | Total 1745 | C 1108 | N 319 | O 314 | S 4 | 0 | 0 | 0 |
| 1 | D | 234 | Total 1786 | C 1133 | N 325 | O 324 | S 4 | 0 | 0 | 0 |
| 1 | E | 226 | Total 1727 | C 1096 | N 317 | O 310 | S 4 | 0 | 0 | 0 |
| 1 | F | 234 | Total 1782 | C 1131 | N 325 | O 322 | S 4 | 0 | 0 | 0 |

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 2 | A | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 2 | A | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 2 | B | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 2 | C | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 2 | D | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 2 | E | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |

- Molecule 3 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|-----|---------|---------|
| 3 | A | 158 | Total | O | 0 | 0 |
| | | | 158 | 158 | | |
| 3 | B | 149 | Total | O | 0 | 0 |
| | | | 149 | 149 | | |
| 3 | C | 113 | Total | O | 0 | 0 |
| | | | 113 | 113 | | |
| 3 | D | 106 | Total | O | 0 | 0 |
| | | | 106 | 106 | | |
| 3 | E | 112 | Total | O | 0 | 0 |
| | | | 112 | 112 | | |
| 3 | F | 128 | Total | O | 0 | 0 |
| | | | 128 | 128 | | |

SEQUENCE-PLOTS INFOmissingINFO

2 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 43 21 2 | Depositor |
| Cell constants a, b, c, α , β , γ | 131.92Å 131.92Å 169.91Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 19.98 – 1.90 19.98 – 1.90 | Depositor EDS |
| % Data completeness (in resolution range) | 98.8 (19.98-1.90) 98.8 (19.98-1.90) | Depositor EDS |
| R_{merge} | 0.06 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 4.09 (at 1.90Å) | Xtrriage |
| Refinement program | CNS 1.1 | Depositor |
| R, R_{free} | 0.180 , 0.208 0.176 , (Not available) | Depositor DCC |
| R_{free} test set | No test flags present. | wwPDB-VP |
| Wilson B-factor (Å ²) | 16.9 | Xtrriage |
| Anisotropy | 0.057 | Xtrriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.42 , 62.9 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$ | Xtrriage |
| Estimated twinning fraction | No twinning to report. | Xtrriage |
| F_o, F_c correlation | 0.95 | EDS |
| Total number of atoms | 11369 | wwPDB-VP |
| Average B, all atoms (Å ²) | 22.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

3 Model quality [i](#)

3.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.3 Torsion angles [i](#)

3.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

3.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

3.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|------|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 2 | GOL | D | 1236 | - | 5,5,5 | 0.29 | 0 | 5,5,5 | 0.17 | 0 |
| 2 | GOL | B | 1236 | - | 5,5,5 | 0.29 | 0 | 5,5,5 | 0.28 | 0 |
| 2 | GOL | A | 1236 | - | 5,5,5 | 0.26 | 0 | 5,5,5 | 0.16 | 0 |
| 2 | GOL | A | 1237 | - | 5,5,5 | 0.21 | 0 | 5,5,5 | 0.12 | 0 |
| 2 | GOL | C | 1236 | - | 5,5,5 | 0.30 | 0 | 5,5,5 | 0.13 | 0 |
| 2 | GOL | E | 1236 | - | 5,5,5 | 0.30 | 0 | 5,5,5 | 0.22 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|------|------|---------|----------|-------|
| 2 | GOL | D | 1236 | - | - | 2/4/4/4 | - |
| 2 | GOL | B | 1236 | - | - | 2/4/4/4 | - |
| 2 | GOL | A | 1236 | - | - | 0/4/4/4 | - |
| 2 | GOL | A | 1237 | - | - | 1/4/4/4 | - |
| 2 | GOL | C | 1236 | - | - | 0/4/4/4 | - |
| 2 | GOL | E | 1236 | - | - | 0/4/4/4 | - |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|------|------|-------------|
| 2 | B | 1236 | GOL | O1-C1-C2-C3 |
| 2 | D | 1236 | GOL | C1-C2-C3-O3 |
| 2 | B | 1236 | GOL | O1-C1-C2-O2 |
| 2 | D | 1236 | GOL | O2-C2-C3-O3 |
| 2 | A | 1237 | GOL | O2-C2-C3-O3 |

There are no ring outliers.

No monomer is involved in short contacts.

3.7 Other polymers [i](#)

There are no such residues in this entry.

3.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

4 Fit of model and data

4.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|-----------------|--------|---------------|-----------------------|-------|
| 1 | A | 228/235 (97%) | -0.24 | 6 (2%) 56 58 | 7, 15, 34, 60 | 0 |
| 1 | B | 234/235 (99%) | -0.17 | 11 (4%) 31 34 | 8, 15, 37, 49 | 0 |
| 1 | C | 228/235 (97%) | -0.03 | 12 (5%) 26 29 | 10, 20, 41, 60 | 0 |
| 1 | D | 234/235 (99%) | 0.31 | 21 (8%) 9 10 | 9, 24, 53, 65 | 0 |
| 1 | E | 226/235 (96%) | -0.12 | 7 (3%) 49 51 | 8, 18, 42, 55 | 0 |
| 1 | F | 234/235 (99%) | 0.04 | 15 (6%) 19 22 | 8, 17, 51, 77 | 0 |
| All | All | 1384/1410 (98%) | -0.03 | 72 (5%) 27 30 | 7, 18, 44, 77 | 0 |

All (72) RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | F | 211 | LEU | 9.0 |
| 1 | F | 209 | PRO | 8.5 |
| 1 | A | 214 | PRO | 7.4 |
| 1 | F | 210 | GLU | 7.2 |
| 1 | F | 207 | GLY | 5.8 |
| 1 | D | 209 | PRO | 5.7 |
| 1 | F | 9 | HIS | 5.4 |
| 1 | F | 208 | ASP | 5.4 |
| 1 | F | 10 | PRO | 5.3 |
| 1 | C | 215 | GLU | 5.2 |
| 1 | E | 205 | ARG | 5.2 |
| 1 | B | 10 | PRO | 4.9 |
| 1 | D | 2 | SER | 4.8 |
| 1 | D | 210 | GLU | 4.6 |
| 1 | A | 213 | PRO | 4.6 |
| 1 | A | 215 | GLU | 4.4 |
| 1 | F | 96 | SER | 4.4 |
| 1 | E | 215 | GLU | 4.3 |
| 1 | F | 216 | VAL | 4.2 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | RSRZ |
|------------|--------------|------------|-------------|-------------|
| 1 | B | 97 | ASP | 4.1 |
| 1 | D | 215 | GLU | 4.0 |
| 1 | F | 213 | PRO | 3.9 |
| 1 | C | 205 | ARG | 3.9 |
| 1 | D | 216 | VAL | 3.8 |
| 1 | D | 207 | GLY | 3.7 |
| 1 | D | 213 | PRO | 3.7 |
| 1 | A | 216 | VAL | 3.7 |
| 1 | C | 2 | SER | 3.6 |
| 1 | D | 211 | LEU | 3.6 |
| 1 | D | 36 | ASN | 3.4 |
| 1 | C | 96 | SER | 3.3 |
| 1 | B | 9 | HIS | 3.3 |
| 1 | D | 205 | ARG | 3.3 |
| 1 | B | 2 | SER | 3.2 |
| 1 | E | 147 | PRO | 3.2 |
| 1 | D | 212 | ALA | 3.2 |
| 1 | F | 2 | SER | 3.2 |
| 1 | D | 35 | GLN | 3.0 |
| 1 | B | 95 | SER | 2.9 |
| 1 | B | 96 | SER | 2.9 |
| 1 | C | 95 | SER | 2.9 |
| 1 | C | 9 | HIS | 2.9 |
| 1 | D | 95 | SER | 2.9 |
| 1 | E | 145 | GLY | 2.9 |
| 1 | B | 214 | PRO | 2.8 |
| 1 | F | 97 | ASP | 2.7 |
| 1 | F | 212 | ALA | 2.7 |
| 1 | F | 95 | SER | 2.7 |
| 1 | F | 215 | GLU | 2.7 |
| 1 | B | 207 | GLY | 2.7 |
| 1 | A | 205 | ARG | 2.6 |
| 1 | C | 216 | VAL | 2.6 |
| 1 | B | 38 | ARG | 2.5 |
| 1 | D | 96 | SER | 2.5 |
| 1 | B | 209 | PRO | 2.5 |
| 1 | E | 100 | PRO | 2.5 |
| 1 | C | 97 | ASP | 2.5 |
| 1 | D | 214 | PRO | 2.4 |
| 1 | C | 212 | PRO | 2.4 |
| 1 | E | 146 | TYR | 2.3 |
| 1 | A | 2 | SER | 2.3 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | D | 206 | ILE | 2.3 |
| 1 | D | 31 | LYS | 2.3 |
| 1 | D | 208 | ASP | 2.3 |
| 1 | C | 100 | PRO | 2.2 |
| 1 | C | 31 | LYS | 2.2 |
| 1 | B | 215 | GLU | 2.2 |
| 1 | E | 2 | SER | 2.2 |
| 1 | C | 36 | ASN | 2.1 |
| 1 | D | 235 | VAL | 2.1 |
| 1 | D | 56 | VAL | 2.1 |
| 1 | D | 99 | ALA | 2.0 |

4.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|------|-------|------|------|----------------------------|-------|
| 2 | GOL | E | 1236 | 6/6 | 0.63 | 0.22 | 47,51,53,55 | 0 |
| 2 | GOL | C | 1236 | 6/6 | 0.64 | 0.23 | 52,54,55,59 | 0 |
| 2 | GOL | D | 1236 | 6/6 | 0.69 | 0.22 | 42,44,45,53 | 0 |
| 2 | GOL | A | 1236 | 6/6 | 0.73 | 0.25 | 43,44,47,53 | 0 |
| 2 | GOL | B | 1236 | 6/6 | 0.77 | 0.21 | 45,46,46,51 | 0 |
| 2 | GOL | A | 1237 | 6/6 | 0.78 | 0.17 | 43,45,46,48 | 0 |

4.5 Other polymers [i](#)

There are no such residues in this entry.