wwPDB X-ray Structure Validation Summary Report (i)

Aug 22, 2023 – 02:25 PM EDT

PDB ID	:	2R6G
Title	:	The Crystal Structure of the E. coli Maltose Transporter
Authors	:	Oldham, M.L.; Khare, D.; Quiocho, F.A.; Davidson, A.L.; Chen, J.
Deposited on	:	2007-09-05
Resolution	:	2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	3%, 79%, 16%
1	B	381	6%, 81%, 15%
2	E	370	3%, 85%, 14%
3	F	514	6%, 74%, 19%, 5%
4	G	296	3%, 79%, 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	C	2	100%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 14713 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose/maltodextrin import ATP-binding protein malK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	371	2876	1819	516	528	13	0	0	0
1	B	372	2882	1822	517	530	13	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	159	GLN	GLU	engineered mutation	UNP Q1R3Q1
A	372	ALA	-	expression tag	UNP Q1R3Q1
A	373	SER	-	expression tag	UNP Q1R3Q1
A	374	ALA	-	expression tag	UNP Q1R3Q1
A	375	SER	-	expression tag	UNP Q1R3Q1
A	376	HIS	-	expression tag	UNP Q1R3Q1
A	377	HIS	-	expression tag	UNP Q1R3Q1
A	378	HIS	-	expression tag	UNP Q1R3Q1
A	379	HIS	-	expression tag	UNP Q1R3Q1
A	380	HIS	-	expression tag	UNP Q1R3Q1
A	381	HIS	-	expression tag	UNP Q1R3Q1
B	159	GLN	GLU	engineered mutation	UNP Q1R3Q1
B	372	ALA	-	expression tag	UNP Q1R3Q1
B	373	SER	-	expression tag	UNP Q1R3Q1
B	374	ALA	-	expression tag	UNP Q1R3Q1
B	375	SER	-	expression tag	UNP Q1R3Q1
B	376	HIS	-	expression tag	UNP Q1R3Q1
B	377	HIS	-	expression tag	UNP Q1R3Q1
B	378	HIS	-	expression tag	UNP Q1R3Q1
B	379	HIS	-	expression tag	UNP Q1R3Q1
B	380	HIS	-	expression tag	UNP Q1R3Q1
B	381	HIS	-	expression tag	UNP Q1R3Q1

- Molecule 2 is a protein called Maltose-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	370	2877	1853	469	549	6	0	0	0

- Molecule 3 is a protein called Maltose transport system permease protein malF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	F	490	3806	2505	606	678	17	0	0	0

- Molecule 4 is a protein called Maltose transport system permease protein malG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	G	284	2187	1468	348	363	8	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
5	C	2	23	12	11	0	0	0

- Molecule 6 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
6	A	1	31	10	5	13	3	0	0
			Total	C	N	O	P		
6	B	1	31	10	5	13	3	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Maltose/maltodextrin import ATP-binding protein malK

Chain A:



• Molecule 1: Maltose/maltodextrin import ATP-binding protein malK

Chain B:



• Molecule 2: Maltose-binding periplasmic protein

Chain E:

• Molecule 3: Maltose transport system permease protein malF

Chain F:



• Molecule 4: Maltose transport system permease protein malG

Chain G:



• Molecule 5: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	73.26Å 95.86Å 109.99Å 87.13° 82.43° 75.75°	Depositor
Resolution (Å)	47.40 – 2.80 35.04 – 2.80	Depositor EDS
% Data completeness (in resolution range)	83.8 (47.40-2.80) 83.8 (35.04-2.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$< I/\sigma(I) >$ [1]	2.58 (at 2.81Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.242 , 0.271 0.240 , 0.269	Depositor DCC
R_{free} test set	1317 reflections (2.02%)	wwPDB-VP
Wilson B-factor (Å^2)	60.1	Xtriage
Anisotropy	0.640	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 37.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	14713	wwPDB-VP
Average B, all atoms (Å^2)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.30	0/2926	0.51	0/3968
1	B	0.30	0/2932	0.49	0/3976
2	E	0.31	0/2946	0.45	0/3998
3	F	0.35	0/3901	0.51	0/5313
4	G	0.35	0/2247	0.49	0/3070
All	All	0.32	0/14952	0.49	0/20325

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2876	0	2944	45	0
1	B	2882	0	2949	41	0
2	E	2877	0	2859	29	0
3	F	3806	0	3836	55	0
4	G	2187	0	2277	34	0
5	C	23	0	21	0	0
6	A	31	0	12	4	0
6	B	31	0	12	0	0
All	All	14713	0	14910	191	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 191 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:SER:HB3	1:A:237:PRO:HD3	1.59	0.82
3:F:414:PHE:HA	3:F:418:THR:HG23	1.60	0.82
1:B:236:SER:HB3	1:B:237:PRO:HD3	1.63	0.80
1:B:260:MET:HB2	1:B:261:PRO:HD2	1.62	0.80
4:G:84:TRP:HB3	4:G:245:LEU:HA	1.68	0.76

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	369/381 (97%)	350 (95%)	13 (4%)	6 (2%)	9 31
1	B	370/381 (97%)	343 (93%)	22 (6%)	5 (1%)	11 34
2	E	368/370 (100%)	350 (95%)	17 (5%)	1 (0%)	41 72
3	F	486/514 (95%)	449 (92%)	31 (6%)	6 (1%)	13 39
4	G	280/296 (95%)	268 (96%)	9 (3%)	3 (1%)	14 41
All	All	1873/1942 (96%)	1760 (94%)	92 (5%)	21 (1%)	14 41

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	21	ASP
1	A	83	SER
1	A	236	SER
1	A	271	GLU
3	F	309	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/323 (97%)	305 (97%)	9 (3%)	42	76
1	B	315/323 (98%)	305 (97%)	10 (3%)	39	73
2	E	297/297 (100%)	287 (97%)	10 (3%)	37	71
3	F	398/424 (94%)	365 (92%)	33 (8%)	11	32
4	G	227/237 (96%)	217 (96%)	10 (4%)	28	61
All	All	1551/1604 (97%)	1479 (95%)	72 (5%)	27	60

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	418	THR
4	G	290	THR
3	F	464	THR
4	G	84	TRP
2	E	138	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	335	GLN
3	F	93	ASN
4	G	282	GLN
2	E	355	GLN
3	F	99	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	GLC	C	1	5	12,12,12	0.54	0	17,17,17	0.77	1 (5%)
5	GLC	C	2	5	11,11,12	0.25	0	15,15,17	0.98	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GLC	C	1	5	-	2/2/22/22	0/1/1/1
5	GLC	C	2	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	C	2	GLC	C1-O5-C5	2.53	115.62	112.19
5	C	1	GLC	C1-O5-C5	2.19	117.79	113.66

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	1	GLC	C4-C5-C6-O6
5	C	1	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	ATP	A	1002	-	26,33,33	0.98	1 (3%)	31,52,52	1.50	6 (19%)
6	ATP	B	1001	-	26,33,33	1.00	1 (3%)	31,52,52	1.46	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ATP	A	1002	-	-	5/18/38/38	0/3/3/3
6	ATP	B	1001	-	-	4/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1002	ATP	C5-C4	2.65	1.47	1.40
6	B	1001	ATP	C5-C4	2.64	1.47	1.40

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	B	1001	ATP	N3-C2-N1	-3.59	123.06	128.68
6	A	1002	ATP	N3-C2-N1	-3.51	123.19	128.68
6	A	1002	ATP	C3'-C2'-C1'	3.27	105.89	100.98
6	A	1002	ATP	PA-O3A-PB	-3.19	121.90	132.83
6	A	1002	ATP	PB-O3B-PG	-2.97	122.63	132.83

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1002	ATP	C5'-O5'-PA-O1A
6	B	1001	ATP	PB-O3A-PA-O5'
6	B	1001	ATP	C5'-O5'-PA-O1A
6	A	1002	ATP	PB-O3A-PA-O5'
6	A	1002	ATP	PB-O3B-PG-O3G

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1002	ATP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ATP A 1002



Ligand ATP B 1001



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	371/381 (97%)	-0.00	11 (2%)	50	40	55, 75, 96, 108	0
1	B	372/381 (97%)	0.16	23 (6%)	20	13	59, 87, 142, 163	0
2	E	370/370 (100%)	-0.08	12 (3%)	47	37	64, 84, 109, 120	0
3	F	490/514 (95%)	0.28	33 (6%)	17	10	60, 84, 123, 144	0
4	G	284/296 (95%)	-0.01	10 (3%)	44	34	55, 70, 102, 130	0
All	All	1887/1942 (97%)	0.08	89 (4%)	31	22	55, 81, 119, 163	0

The worst 5 of 89 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	59	ASN	11.7
2	E	170	LYS	5.1
1	B	293	SER	5.1
1	A	372	ALA	5.1
3	F	40	TYR	4.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	GLC	C	1	12/12	0.93	0.25	77,77,78,78	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	GLC	C	2	11/12	0.95	0.34	76,76,76,77	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	ATP	A	1002	31/31	0.94	0.17	58,64,65,65	0
6	ATP	B	1001	31/31	0.95	0.15	63,66,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ATP A 1002:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around ATP B 1001:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.