



wwPDB EM Validation Summary Report ⓘ

May 7, 2024 – 02:47 pm BST

PDB ID : 8R6U
EMDB ID : EMD-18963
Title : Structure of the SFTSV L protein in a transcription-priming state without capped RNA [TRANSCRIPTION-PRIMING (in vitro)]
Authors : Williams, H.M.; Thorkelsson, S.R.; Vogel, D.; Busch, C.; Milewski, M.; Cusack, S.; Grunewald, K.; Queminn, E.R.J.; Rosenthal, M.
Deposited on : 2023-11-23
Resolution : 2.98 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2084	
2	E	20	
2	P	20	
3	T	26	
4	G	16	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase L.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1566	Total	C	N	O	S	2	0
			12495	7936	2170	2317	72		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	112	ALA	ASP	engineered mutation	UNP U3GU88

- Molecule 2 is a RNA chain called RNA primer.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	P	16	Total	C	N	O	P	0	0
			341	154	68	104	15		
2	E	6	Total	C	N	O	P	0	0
			129	58	26	39	6		

- Molecule 3 is a RNA chain called RNA (5'-R(P*CP*UP*GP*GP*GP*CP*GP*GP*UP*CP*UP*UP*UP*GP*UP*GP*U)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
3	T	17	Total	C	N	O	P	0	0
			361	160	58	126	17		

- Molecule 4 is a RNA chain called RNA primer tail-end.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	4	Total	C	N	O	P	0	0
			81	38	16	24	3		


- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
5	A	1	Total	Mg	0
			1	1	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase L



ARG	THR	LEU	ALA	LEU	THR	PRO
GLY	LEU	VAL	TRP	ASN	LEU	GLY
LEU	MET	ASP	SER	ASP	TRP	ILE
ASP	ASP	ILE	HIS	SER	ASN	ASN
GLU	GLN	LEU	ALA	VAL	VAL	ARG
GLU	ALA	GLU	SER	THR	THR	VAL
SER	ALA	GLU	VAL	ILE	ILE	GLY
THR	ILE	ASP	LEU	LEU	GLN	LYS
ILE	ILE	ILE	LEU	GLU	GLU	GLY
MET	MET	ASP	ALA	GLY	GLY	LYS
MET	MET	PHE	ASN	ARG	ARG	GLY
TRP	CYS	ASP	ASP	VAL	VAL	GLY
GLY	ALA	SER	ASP	VAL	VAL	VAL
	ILE	ASP	ARG	MET	MET	TRP
	ILE	ASP	ARG	MET	HIS	TRP
	MET	VAL	LYS	ASN	CYS	THR
	GLY	ILE	THR	ILE	MET	GLY
	LYS	ALA	GLN	LEU	VAL	VAL
	GLU	VAL	GLY	SER	ARG	VAL
	GLY	GLU	ILE	TYR	LEU	GLU
	CYS	LEU	ASP	TYR	SER	ASP
	ARG	SER	ASN	PRO	GLY	THR
	GLY	GLU	ARG	ARG	PHE	HIS
	LEU	GLY	ALA	ASP	LYS	VAL
	LEU	SER	MET	THR	ILE	GLN
	THR	LEU	GLY	ASP	LYS	ILE
	GLU	ASP	ASN	ILE	PRO	LEU
	LYS	ILE	ILE	SER	ALA	ILE
	ARG	GLU	PHE	GLU	SER	ASP
	CYS	SER	ARG	SER	GLY	GLY
	MET	ILE	ASP	ALA	THR	ASP
	ALA	PHE	CYS	ALA	ASP	GLY
	ALA	ASP	LEU	ALA	GLY	THR
	ILE	GLY	LEU	TYR	CYS	SER
	ARG	ALA	GLY	LEU	PRO	ASN
	GLU	PRO	SER	TRP	VAL	TRP
	GLN	ILE	LEU	SER	ARG	LEU
	VAL	LEU	ARG	ASN	ILE	GLU
	ARG	TRP	LYS	ARG	MET	GLU
	PRO	SER	GLN	ASP	GLU	ILE
	PHE	ALA	GLY	LEU	ARG	ARG
	LEU	GLU	LEU	PHE	GLY	LEU
	ILE	VAL	MET	SER	PHE	SER
	PHE	GLU	ARG	PHE	ARG	SER
	LEU	GLU	SER	GLY	ILE	ASP
	GLN	PHE	LYS	LYS	ARG	ALA
	PRO	ILE	LEU	LEU	GLU	ARG
	GLU	GLU	THR	GLU	LEU	LEU
	GLY	GLY	GLU	PRO	GLN	TYR
	VAL	VAL	MET	SER	ASN	ASP
	SER	VAL	VAL	CYS	ASP	ASP
	SER	SER	VAL	LEU	VAL	VAL
	ASP	SER	PRO	LYS	ARG	ARG
	GLN	LYS	LEU	THR	VAL	ARG
	PHE	THR	THR	LEU	GLY	CYS
	CYS	THR	THR	ASP	GLY	ASP
	ASP	HIS	GLN	ASN	ASP	ASP
	SER	LEU	THR	THR	ILE	ILE

- Molecule 2: RNA primer


Chain P:  55% 20% 5% 20%

- Molecule 2: RNA primer

Chain E: 20% 5% 5% 70%

A C2 A3 A7 G A C C G C C C A G A U G A

- Molecule 3: RNA (5'-R(P*CP*UP*GP*GP*GP*CP*GP*GP*UP*CP*UP*UP*UP*GP*UP*GP*U)-3')

Chain T:  27% 19% 19% 35%

A	A	A	A	A	A	A	G	A	U	C10	U11	G12	G13	G14	C15	G16	G17	U18	C19	U20	U21	U22		U26
---	---	---	---	---	---	---	---	---	---	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	--	-----

- Molecule 4: RNA primer tail-end

Chain G:  12% 6% 6% 75%

A	A	A	A	C	G	C	A	A	C	C	A	A1	C2	A3	C4
---	---	---	---	---	---	---	---	---	---	---	---	----	----	----	----

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	67479	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.036	Depositor
Minimum map value	-0.018	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.004	Depositor
Map size (Å)	272.0, 272.0, 272.0	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.85, 0.85, 0.85	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/12766	0.54	0/17228
2	E	0.40	0/144	0.73	0/222
2	P	0.74	0/382	0.70	0/594
3	T	0.47	0/401	0.93	1/623 (0.2%)
4	G	0.56	0/90	1.62	1/138 (0.7%)
All	All	0.40	0/13783	0.58	2/18805 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	4	C	OP1-P-OP2	-6.49	109.87	119.60
3	T	22	U	N1-C2-O2	5.15	126.40	122.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	12495	0	12513	124	0
2	E	129	0	67	1	0
2	P	341	0	178	1	0
3	T	361	0	181	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	81	0	46	0	0
5	A	1	0	0	0	0
All	All	13408	0	12985	127	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 127 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:ILE:HG13	1:A:956:ARG:HD3	1.77	0.67
1:A:121:THR:HG22	1:A:164:ARG:HB2	1.76	0.67
1:A:345:ASP:O	1:A:571:ARG:NH1	2.28	0.67
1:A:915:GLN:HG3	1:A:917:GLY:H	1.60	0.66
1:A:926:ASP:OD1	1:A:926:ASP:N	2.29	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1558/2084 (75%)	1455 (93%)	103 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1378/1826 (76%)	1296 (94%)	82 (6%)	19	51

5 of 82 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1290	ARG
1	A	1452	HIS
1	A	1339	MET
1	A	1375	ARG
1	A	1507	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	964	ASN
1	A	981	ASN
1	A	1599	GLN
1	A	1353	ASN
1	A	1467	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	E	5/20 (25%)	2 (40%)	0
2	P	15/20 (75%)	4 (26%)	0
3	T	16/26 (61%)	6 (37%)	0
4	G	3/16 (18%)	2 (66%)	0
All	All	39/82 (47%)	14 (35%)	0

5 of 14 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	P	5	A
2	P	7	A
2	P	11	G
2	P	16	G
3	T	11	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-18963. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 157




Y Index: 153




Z Index: 137


6.3.2 Raw map



X Index: 158



Y Index: 157




Z Index: 147


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.004. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 138 nm³; this corresponds to an approximate mass of 125 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.336 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.336 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.98	-	-
Author-provided FSC curve	2.96	3.42	2.99
Unmasked-calculated*	3.76	6.68	3.89

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.76 differs from the reported value 2.98 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-18963 and PDB model 8R6U. Per-residue inclusion information can be found in section [3](#) on page [5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.004 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.004).

9.4 Atom inclusion [i](#)



At the recommended contour level, 96% of all backbone atoms, 93% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.004) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.9340	<div></div> 0.5650
A	<div></div> 0.9360	<div></div> 0.5690
E	<div></div> 0.8910	<div></div> 0.4750
G	<div></div> 0.6910	<div></div> 0.3170
P	<div></div> 0.9820	<div></div> 0.5820
T	<div></div> 0.8730	<div></div> 0.4890

