



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2023 – 06:23 AM EDT

PDB ID : 6UBH
Title : Structure of the MM7 Erbin PDZ variant in complex with a high-affinity peptide
Authors : Singer, A.U.; Teyra, J.; McLaughlin, M.; Ernst, A.; Sicheri, F.; Sidhu, S.S.
Deposited on : 2019-09-11
Resolution : 1.80 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3333 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Erbin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	91	694	438	121	133	2	0	0	0
1	B	92	693	437	122	132	2	0	0	0
1	C	92	717	452	125	136	4	8	2	0
1	D	92	709	447	124	135	3	0	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	SER	-	expression tag	UNP Q96RT1
A	20	SER	-	expression tag	UNP Q96RT1
A	21	MET	-	expression tag	UNP Q96RT1
A	89	MET	HIS	engineered mutation	UNP Q96RT1
B	19	SER	-	expression tag	UNP Q96RT1
B	20	SER	-	expression tag	UNP Q96RT1
B	21	MET	-	expression tag	UNP Q96RT1
B	89	MET	HIS	engineered mutation	UNP Q96RT1
C	19	SER	-	expression tag	UNP Q96RT1
C	20	SER	-	expression tag	UNP Q96RT1
C	21	MET	-	expression tag	UNP Q96RT1
C	89	MET	HIS	engineered mutation	UNP Q96RT1
D	19	SER	-	expression tag	UNP Q96RT1
D	20	SER	-	expression tag	UNP Q96RT1
D	21	MET	-	expression tag	UNP Q96RT1
D	89	MET	HIS	engineered mutation	UNP Q96RT1

- Molecule 2 is a protein called peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	5	Total	C	N	O	0	0	0
			46	32	6	8			
2	F	5	Total	C	N	O	0	0	0
			46	32	6	8			
2	G	5	Total	C	N	O	0	0	0
			46	32	6	8			
2	H	5	Total	C	N	O	0	0	0
			52	38	6	8			

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		
3	D	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	83	Total	O	0	0
			83	83		
4	B	74	Total	O	0	0
			74	74		
4	C	78	Total	O	0	0
			78	78		
4	D	79	Total	O	0	0
			79	79		
4	E	2	Total	O	0	0
			2	2		
4	F	6	Total	O	0	0
			6	6		
4	G	3	Total	O	0	0
			3	3		
4	H	2	Total	O	0	0
			2	2		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	39.00Å 39.01Å 58.38Å 72.55° 72.61° 96.06°	Depositor
Resolution (Å)	36.77 – 1.80	Depositor
% Data completeness (in resolution range)	81.5 (36.77-1.80)	Depositor
R_{merge}	0.02	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	15.27 (at 1.81Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.197 , 0.225	Depositor
Wilson B-factor (Å ²)	16.2	Xtrriage
Anisotropy	0.048	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	0.479 for k,h,h+k-l 0.460 for -k,-h,-l 0.466 for -h,-k,-h-k+l	Xtrriage
Total number of atoms	3333	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.